

Strong Duality and Minimal Representations for Cone Optimization

Levent Tunçel Henry Wolkowicz

August 2008, revised: February 23, 2011

University of Waterloo
Department of Combinatorics & Optimization
Waterloo, Ontario N2L 3G1, Canada
Research Report CORR 2008-07

Key words and phrases: Cone Optimization, Duality, Preprocessing, Constraint Qualification, Duality Gap, Semidefinite Programming, Strict Complementarity, Nice Cones, Devious Cones, Facially Dual Complete Cones.

AMS subject classifications: 90C46, 90C22, 49K40, 90C31

Abstract

The elegant theoretical results for strong duality and strict complementarity for linear programming, LP, lie behind the success of current algorithms. In addition, preprocessing is an essential step for efficiency in both simplex type and interior-point methods. However, the theory and preprocessing techniques can fail for cone programming over nonpolyhedral cones. We take a fresh look at known and new results for duality, optimality, constraint qualifications, CQ, and strict complementarity, for linear cone optimization problems in finite dimensions. One theme is the notion of *minimal representation* of the cone and the constraints. This provides a framework for preprocessing cone optimization problems in order to avoid both the theoretical and numerical difficulties that arise due to the (near) loss of the strong CQ, strict feasibility. We include results and examples on the surprising theoretical connection between duality gaps in the original primal-dual pair and lack of strict complementarity in their homogeneous counterpart. Our emphasis is on results that deal with Semidefinite Programming, SDP.

Contents

1	Introduction	2
1.1	Background and Motivation	3
1.2	Outline	3
2	Notation and Preliminary Results	4
2.1	Subspace Form, Complementarity Partitions, and Minimal Sets	5
2.1.1	Subspace Form for Primal-Dual Pair (\mathbb{P}) and (\mathbb{D})	5
2.1.2	Complementarity Partitions, CP, and Minimal Sets	6

3	Facial Properties	8
3.1	Faces of General Cones	9
3.2	Faces for Primal-Dual Pair (\mathbb{P}) and (\mathbb{D})	10
3.3	Nice Cones, Devious Cones, and SDP	12
4	Duality and Minimal Representations	14
4.1	Strong Duality and Constraint Qualifications	14
4.1.1	Universal Constraint Qualifications	18
5	Failure of Strong Duality and Strict Complementarity	18
5.1	Finite Positive Duality Gaps	19
5.1.1	Positive Gaps and Closure Conditions	19
5.1.2	Positive Gaps and Strict Complementarity	20
5.2	Infinite Duality Gap and Devious Faces	25
5.3	Regularization for Strict Complementarity	26
6	Conclusion	26
	Bibliography	26
	Index	32
A	comments from referee	33

1 Introduction

In this paper we study duality, optimality conditions, and preprocessing for the *conic optimization problem*, i.e., the problem of optimizing a linear objective function over the intersection of a convex cone with an affine space. We include both the linear transformation and the subspace forms for our optimization problems; we study known and new characterizations of optimality that hold without any constraint qualification, CQ; and, we collect needed technical results on the cone facial structure.

In addition, we derive new CQs including a *universal* constraint qualification, UCQ, i.e., a CQ that holds independent of the data b, c ; and, we study the geometry of *nice* and *devious* cones and the relationship that lack of closure has to both strong duality and zero duality gaps, including characterizations for a zero duality gap, and a surprising connection between duality gaps and the failure of strict complementarity in the homogeneous problems.

One theme is the notion of *minimal representation* of the cone and the constraints in order to preprocess and regularize the problem and thus avoid both the theoretical and numerical difficulties that arise due to (near) loss of strict feasibility, i.e., we see that loss of this strong CQ is a modelling issue rather than inherent to the problem instance. As it is well-known that the existence of a strong CQ of the Mangasarian-Fromovitz or Robinson type is equivalent to stability of the problem, e.g., [65], this justifies the pleasing paradigm: *efficient modelling provides for a stable program*.

1.1 Background and Motivation

The cone programming model has been studied for a long time, e.g., [23], as a generalization of the classical linear program, LP. More recently, many important applications have arisen for more general nonpolyhedral cones. This includes the case when K is the cone of positive semidefinite matrices, \mathbb{S}_+^n ; then we get semidefinite programming, SDP, e.g., [73]. Another important case is second order cone programming, SOCP, where $K = \text{SOC}_1 \oplus \cdots \oplus \text{SOC}_k$, a direct sum of second order (Lorentz) cones, e.g., [4, 45]. These research areas remain very active, see e.g., [2, 33, 38, 41, 49, 67, 73, 77] and URL: www-user.tu-chemnitz.de/~helmberg/semidef.html. Optimality conditions and CQs have been studied in e.g., [23, 31, 62] and more recently for both linear and nonlinear problems in e.g., [65]. (See the historical notes in [65, Sect. 4.1.5].) Optimality conditions and strong duality results without a CQ have appeared in e.g., [13, 14, 15, 16, 35, 36, 53, 57, 58].

Both strong duality and strict complementarity behave differently for general cone optimization problems, compared to the LP case. First, strong duality for a cone program can fail in the absence of a CQ, i.e., there may not exist a dual optimal solution and there may be a nonzero duality gap. In addition, the (near) failure of the Slater CQ (strict feasibility) has been used in complexity measures, [60, 61]. Moreover, numerical difficulties are well correlated with (near) failure of the Slater CQ, see [25, 26]. Similarly, unlike the LP case, [28], there are general cone optimization problems for which there does not exist a primal-dual optimal solution that satisfies strict complementarity, see e.g., [73] for examples. Theoretical difficulties arise, e.g., for local convergence rate analysis. Again, we have that numerical difficulties are well correlated with loss of strict complementarity, see [71]. An algorithm for generating SDP problems where strict complementarity fails, independent of whether the Slater CQ holds or not, is also given in [71].

Connections between weakest CQs and the closure of the sum of a subspace and a cone date back to e.g., [31]. We present a surprising theoretical connection between strict complementarity of the homogeneous problem and duality gaps, as well as show that both loss of strict complementarity and strong duality are connected to the lack of closure of the sum of a cone and a subspace.

Examples where no CQ holds arise in surprisingly many cases. For example, Slater's CQ fails for many SDP relaxations of hard combinatorial problems, see e.g., [5, 74, 75]. A unifying approach to remedy this situation is given in [68]. Another instance is the SDP that arises from relaxations of polynomial optimization problems, e.g., [69]. Exploiting the absence of Slater's CQ is done in [42]. Current public domain codes for SDP are based on interior-point methods and do not take into account loss of Slater's CQ (strict feasibility) or loss of strict complementarity. As discussed above, both of these conditions can result in theoretical and numerical problems, e.g., [25, 69, 71]. Contrary to the LP case, e.g., [29, 39, 48], current SDP codes do not perform extensive preprocessing to avoid these difficulties. (Though some preprocessing is done to take advantage of sparsity, e.g., [27]. A projection technique for the cases where Slater's CQ fails is studied in [18].)

1.2 Outline

In Section 2 we present the notation and preliminary results. We introduce: the subspace forms for the cone optimization in Section 2.1.1 and the complementarity partition and minimal sets in Section 2.1.2. Technical facial properties are presented in Sections 3.1 and 3.2. The notions of nice and devious cones are described in Section 3.3. We include many relationships for the facial structure of the cone optimization problems.

The strong duality results, with and without CQs, and the CQs and UCQ, are presented in

Section 4, see e.g., Theorem 4.10. We use both the minimal cone known in the literature and introduce the minimal subspace in order to obtain a regularization that guarantees that Slater's CQ holds, rather than the weaker generalized Slater CQ given in the literature, see (4.9) in Theorem 4.10. We study the failure of strong duality and strict complementarity in Section 5. This includes a characterization for a zero duality gap in Section 5.1. The surprising relation between duality gaps and the failure of the strict complementarity property for the homogeneous problem, is given in Section 5.1.2, see e.g., Theorems 5.12 and 5.10. Our concluding remarks are in Section 6.

2 Notation and Preliminary Results

The set K is a convex cone if it is a cone, i.e., it is closed under nonnegative scalar multiplication, $\lambda K \subseteq K, \forall \lambda \geq 0$, and, it is also closed under addition $K + K \subseteq K$. The cone K is a proper cone if it is closed, pointed, and has nonempty interior. We let $u \preceq_K v$ (respectively, $u \prec_K v$) denote the partial order induced by K , i.e., $v - u \in K$ (respectively, $v - u \in \text{int } K$).

We use \bar{S} to denote *closure*, $\text{precl } S = \bar{S} \setminus S$ to denote the *preclosure* of a set S . We let $\text{conv } S$ denote the *convex hull of the set* S and $\text{cone } S$ denote the *convex cone generated by* S . (By abuse of notation, we use $\text{cone } s := \text{cone } \{s\}$, for a single element s . This holds similarly for, e.g., $s^\perp := \{s\}^\perp$ and other operations that act on single element sets.) The dual or nonnegative polar cone of a set S is $S^* := \{x : \langle x, s \rangle \geq 0, \forall s \in S\}$. In particular, for the space of $n \times n$ symmetric matrices, \mathbb{S}^n , we use the trace inner-product $\langle x, s \rangle = \text{trace } xs$, i.e., the trace of the product of the matrices x and s . We use $\mathcal{R}(\cdot)$ and $\mathcal{N}(\cdot)$ to denote range space and nullspace, respectively. We let $\mathbb{S}_+^n \subset \mathbb{S}^n$ denote the cone of positive semidefinite matrices. In particular, setting $K = \mathbb{S}_+^n$ yields the partial order induced by the cone of positive semidefinite matrices in \mathbb{S}^n , i.e., the so-called Löwner partial order, $x \preceq_{\mathbb{S}_+^n} s$.

We let e_i denote the i th unit vector of appropriate dimension, and E_{ij} denote the (i, j) th unit matrix in \mathbb{S}^n , i.e., $E_{ii} = e_i e_i^T$ and if $i \neq j$, $E_{ij} = e_i e_j^T + e_j e_i^T$. By abuse of notation, we let x_{ij} denote the ij element of $x \in \mathbb{S}^n$.

The subset $F \subseteq K$ is a *face of the cone* K , denoted $F \trianglelefteq K$, if

$$(s \in F, 0 \preceq_K u \preceq_K s) \text{ implies } (\text{cone } u \subseteq F). \quad (2.1)$$

If $F \trianglelefteq K$ but is not equal to K , we write $F \triangleleft K$. If $\{0\} \subset F \triangleleft K$, then F is a *proper face* of K . (Similarly, $S_1 \subset S_2$ denotes a proper subset, i.e., $S_1 \subseteq S_2, S_1 \neq S_2$.) For $S \subseteq K$, we let $\text{face } S$ denote the smallest face of K that contains S ; equivalently $\text{face } S$ is the intersection of all faces containing S . A face $F \trianglelefteq K$ is an *exposed face* if it is the intersection of K with a hyperplane. The cone K is *facially exposed* if every face $F \trianglelefteq K$ is exposed. If $F \trianglelefteq K$, then the *conjugate face* is $F^c := K^* \cap F^\perp$. Note that if the conjugate face F^c is a proper face, then it is *exposed* using any $s \in \text{relint } F$, i.e., $F^c = K \cap s^\perp, \forall s \in \text{relint } F$.

We study the following pair of dual conic optimization problems in standard form:

$$(\mathbb{P}) \quad v_P := \sup_y \{ \langle b, y \rangle : \mathcal{A}^* y \preceq_K c \} \quad (2.2)$$

$$(\mathbb{D}) \quad v_D := \inf_x \{ \langle c, x \rangle : \mathcal{A} x = b, x \succeq_{K^*} 0 \}, \quad (2.3)$$

where: $\mathcal{A} : \mathcal{V} \rightarrow \mathcal{W}$ is a (onto) linear transformation between two finite dimensional inner-product spaces; \mathcal{A}^* denotes the adjoint transformation; and K is a convex cone. Throughout, we assume

that the optimal value v_P is finite. *Weak duality* holds for any pair of primal-dual feasible solutions y, x , i.e., if $s = c - \mathcal{A}^* y \succeq_K 0$, $\mathcal{A}x = b$, $x \succeq_{K^*} 0$, then we get

$$\langle b, y \rangle = \langle \mathcal{A}x, y \rangle = \langle \mathcal{A}^* y, x \rangle = \langle c - s, x \rangle \leq \langle c, x \rangle. \quad (\text{Weak Duality})$$

The usual constraint qualification, CQ, used for the primal (2.2) is the Slater condition, i.e., strict feasibility $\mathcal{A}^* \hat{y} \prec c$. If we assume Slater's CQ holds and the primal optimal value is finite, then strong duality holds, i.e., we have a zero duality gap and attainment of the dual optimal value,

$$v_P = v_D = \langle c, x^* \rangle, \text{ for some dual feasible } x^*. \quad (\text{Strong Duality})$$

Denote the primal-dual feasible sets of (2.2) and (2.3) by

$$\mathcal{F}_P^y = \mathcal{F}_P^y(c) = \{y : \mathcal{A}^* y \preceq_K c\}, \quad \mathcal{F}_D^x = \mathcal{F}_D^x(b) = \{x : \mathcal{A}x = b, x \succeq_{K^*} 0\}, \quad (2.4)$$

respectively. The set of feasible slacks for (2.2) is

$$\mathcal{F}_P^s = \mathcal{F}_P^s(c) = \{s : s = c - \mathcal{A}^* y \succeq_K 0, \text{ for some } y\}. \quad (2.5)$$

We allow for the dependence on the parameters b and c . Similarly, the *optimal solution sets* are denoted by $\mathcal{O}_P^s, \mathcal{O}_P^y, \mathcal{O}_D^x$. Moreover, the pair of feasible primal-dual solutions s, x are said to satisfy *strict complementarity, SC* if

$$\begin{aligned} (\text{SC}) \quad & s \in \text{relint } F_P \text{ and } x \in \text{relint } F_P^c, \quad \text{for some } F_P \trianglelefteq K, \\ & \text{or} \\ & s \in \text{relint } F_D^c \text{ and } x \in \text{relint } F_D, \quad \text{for some } F_D \trianglelefteq K^*. \end{aligned} \quad (2.6)$$

(Note that this implies $s + x \in \text{int}(K + K^*)$, see Proposition 3.3 part 1, below.)

2.1 Subspace Form, Complementarity Partitions, and Minimal Sets

2.1.1 Subspace Form for Primal-Dual Pair (P) and (D)

Suppose that \tilde{s}, \tilde{y} , and \tilde{x} satisfy

$$\mathcal{A}^* \tilde{y} + \tilde{s} = c, \quad \mathcal{A} \tilde{x} = b. \quad (2.7)$$

Then, for any feasible primal-dual triple (x, y, s) , where s is, as usual, the primal slack given by $s = c - \mathcal{A}^* y$, we have $\langle c, \tilde{x} \rangle = \langle \mathcal{A}^* y + s, \tilde{x} \rangle = \langle b, y \rangle + \langle s, \tilde{x} \rangle$. Therefore, the objective in (2.2) can be rewritten as

$$\sup_y \langle b, y \rangle = \sup_s (\langle c, \tilde{x} \rangle - \langle s, \tilde{x} \rangle) = \langle c, \tilde{x} \rangle - \inf_s \langle s, \tilde{x} \rangle.$$

We let \mathcal{L} denote the nullspace $\mathcal{N}(\mathcal{A})$ of the operator \mathcal{A} . Then

$$\mathcal{F}_P^s = \mathcal{F}_P^s(c) = (c + \mathcal{L}^\perp) \cap K = (\tilde{s} + \mathcal{L}^\perp) \cap K. \quad (2.8)$$

In addition, for $x \in \tilde{x} + \mathcal{L}$, we get $\langle c, x \rangle = \langle \mathcal{A}^* \tilde{y} + \tilde{s}, x \rangle = \langle \tilde{s}, x \rangle + \langle \mathcal{A}^* \tilde{y}, x \rangle = \langle \tilde{s}, x \rangle + \langle \tilde{y}, b \rangle$. We can now write the primal and dual conic pair, (2.2) and (2.3), in the so-called subspace form (see e.g., [50, Section 4.1]):

$$v_P = \langle c, \tilde{x} \rangle - \inf_s \{\langle s, \tilde{x} \rangle : s \in (\tilde{s} + \mathcal{L}^\perp) \cap K\} \quad (2.9)$$

$$v_D = \langle \tilde{y}, b \rangle + \inf_x \{ \langle \tilde{s}, x \rangle : x \in (\tilde{x} + \mathcal{L}) \cap K^* \}. \quad (2.10)$$

The symmetry means that we can directly extend results proved for (2.9) to (2.10). Note that we have much flexibility in the choice of \tilde{s} and \tilde{x} . In particular, if (2.9) and (2.10) are feasible, we may choose $\tilde{s} \in \mathcal{F}_P^s$ and $\tilde{x} \in \mathcal{F}_D^x$, and in the case that the optimal values are attained, we may choose $\tilde{s} \in \mathcal{O}_P^s$ and $\tilde{x} \in \mathcal{O}_D^x$.

Proposition 2.1 *Let \tilde{s}, \tilde{y} , and \tilde{x} satisfy (2.7). Then (2.9) and (2.10) are a dual pair of cone optimization problems equivalent to (2.2) and (2.3), respectively. Moreover, (2.2) (resp. (2.3)) is feasible if and only if $\tilde{s} \in K + \mathcal{L}^\perp$ (resp. $\tilde{x} \in K^* + \mathcal{L}$). ■*

2.1.2 Complementarity Partitions, CP, and Minimal Sets

Denote the minimal faces for the homogeneous problems (recession directions) by

$$f_P^0 := \text{face } \mathcal{F}_P^s(0) = \text{face } (\mathcal{L}^\perp \cap K) \quad (2.11)$$

$$f_D^0 := \text{face } \mathcal{F}_D^x(0) = \text{face } (\mathcal{L} \cap K^*). \quad (2.12)$$

For connections between recession directions and optimality conditions, see e.g., [1, 6, 11]. Note that $f_P^0 \subseteq (f_D^0)^c$ (equivalently, $f_D^0 \subseteq (f_P^0)^c$).

Definition 2.2 *The pair of faces $F_1 \trianglelefteq K, F_2 \trianglelefteq K^*$ form a complementarity partition of K, K^* , denoted CP, if $F_1 \subseteq F_2^c$. (Equivalently, $F_2 \subseteq F_1^c$.) The partition is proper if both F_1 and F_2 are proper faces. The partition is strict if $(F_1)^c = F_2$ or $(F_2)^c = F_1$.*

It is well known that

$$F \trianglelefteq K, G \trianglelefteq K, F \subseteq G \implies F \trianglelefteq G. \quad (2.13)$$

Therefore, we can assume $F_1 \trianglelefteq F_2^c$ and $F_2 \trianglelefteq F_1^c$ in Definition 2.2. Moreover, for every linear subspace \mathcal{L} , the pair of faces

$$(f_P^0, f_D^0) \text{ form a complementarity partition, CP} \quad (2.14)$$

of K, K^* ; and, if K is a polyhedral cone, then the partition is strict. The *minimal face* of (2.2) is the face of K generated by the feasible slack vectors; while the *minimal face* for (2.3) is the face of K^* generated by the feasible set of the dual problem, i.e., we denote $f_P := \text{face } \mathcal{F}_P^s$, $f_D := \text{face } \mathcal{F}_D^x$. Note that both $f_P = f_P(\tilde{s})$ and $f_D = f_D(\tilde{x})$, i.e., they depend implicitly on the points \tilde{s}, \tilde{x} in the subspace formulations (2.9) and (2.10).

An immediate important property of the complementarity partitions is:

$$\text{if } (F_1, F_2) \text{ form a CP, then } \boxed{0 \neq \phi \in F_1 \implies F_2 \subseteq \{\phi\}^\perp \implies \text{int } F_2 = \emptyset} \quad (2.15)$$

We now obtain the following complementarity partitions and the corresponding theorems of the alternative from (2.15).

Proposition 2.3 *Let $\tilde{y}, \tilde{s}, \tilde{x}$ satisfy (2.7). Then the five pairs of faces*

$$\left[\begin{array}{cc} f_P^0 & f_D^0 \\ \text{face } ((\mathcal{L}^\perp + \text{span } \tilde{s}) \cap K) & \text{face } (f_D^0 \cap \tilde{s}^\perp) \\ \text{face } (f_P^0 \cap \tilde{x}^\perp) & \text{face } ((\mathcal{L} + \text{span } \tilde{x}) \cap K^*) \\ f_P & \text{face } (f_D^0 \cap \tilde{s}^\perp) \\ \text{face } (f_P^0 \cap \tilde{x}^\perp) & f_D \end{array} \right] \quad (2.16)$$

form complementarity partitions of K, K^* .

Proof.

That the first pair in (2.16) forms a complementarity partition follows from definitions. The result for the second and third pairs follow from replacing \mathcal{L}^\perp with $\mathcal{L}^\perp + \text{span } \tilde{s}$, and from replacing \mathcal{L} with $\mathcal{L} + \text{span } \tilde{x}$, respectively. The result for the final two pairs follows from

$$\mathcal{F}_P^s \subseteq \left(\mathcal{L}^\perp + \text{span } \tilde{s} \right) \cap K, \quad \mathcal{F}_D^x \subseteq \left(\mathcal{L} + \text{span } \tilde{x} \right) \cap K^*.$$

■

The primal and dual minimal subspace representations of \mathcal{L}^\perp and of \mathcal{L} , respectively, are given by

$$\mathcal{L}_{PM}^\perp := \mathcal{L}^\perp \cap (f_P - f_P), \quad \mathcal{L}_{DM} := \mathcal{L} \cap (f_D - f_D). \quad (2.17)$$

The cone of feasible directions at $\hat{y} \in \mathcal{F}_P^y$ is

$$\mathcal{D}_P^{\leq}(\hat{y}) = \text{cone} \left(\mathcal{F}_P^y - \hat{y} \right). \quad (2.18)$$

We similarly define the cones $\mathcal{D}_P^{\leq}(\hat{s}), \mathcal{D}_D^{\leq}(\hat{x})$. For these three cones, we assume that $\hat{y}, \hat{s}, \hat{x}$ are suitable feasible points in $\mathcal{F}_P^y, \mathcal{F}_P^s, \mathcal{F}_D^x$, respectively. The closures of these cones of feasible directions yield the standard *tangent cones*, denoted $\mathbb{T}_P(\hat{y}), \mathbb{T}_P(\hat{s}), \mathbb{T}_D(\hat{x})$, respectively. (See e.g., [9, 21].) Note that if the primal feasible set is simply K , the cone of feasible directions corresponds to the so-called *radial cone*.

Proposition 2.4 ([66, 70]) *Let K be closed. Then K is a polyhedral cone if and only if at every point $\hat{s} \in K$, the radial cone of K , $\text{cone}(K - \hat{s})$, at \hat{s} is closed.* ■

Example 2.5 *We now look at two examples that illustrate the lack of closure for nonpolyhedral cones, e.g., in each instance we get*

$$K + \text{span } f_P^0 \subset \overline{K + \text{span } f_P^0} = K + ((f_P^0)^c)^\perp. \quad (2.19)$$

The lack of closure in (2.19) can be used to find examples with both finite and infinite positive duality gaps; see Theorem 5.3 and the following example in item 1.

1. First, let $n = 2$ and \mathcal{L} in (2.9) and (2.10) be such that $\mathcal{L}^\perp = \text{span}\{E_{22}\}$. Then $f_P^0 = \text{cone}\{E_{22}\}$ and $f_D^0 = \text{cone}\{E_{11}\}$. Therefore, $f_P^0 = (f_D^0)^c$ and $f_D^0 = (f_P^0)^c$, i.e., this is a strict complementarity partition. Moreover, (2.19) holds e.g., $E_{12} \in (f_P^0)^\perp \cap (f_D^0)^\perp$ and

$$E_{12} = \lim_{i \rightarrow \infty} \left(\begin{bmatrix} 1/i & 1 \\ 1 & i \end{bmatrix} - iE_{22} \right) \in \left(\mathbb{S}_+^2 + (f_D^0)^\perp \right) \setminus \left(\mathbb{S}_+^2 + \text{span } f_P^0 \right) = \text{precl} \left(\mathbb{S}_+^2 + \text{span } f_P^0 \right).$$

We can now choose $\tilde{s} = E_{12}, \tilde{x} = E_{11}$. Then the primal is infeasible, e.g., Proposition 2.1, while the dual optimal value $v_D = 0$.

2. Now, let $n = 3$ and suppose that $\mathcal{L}^\perp = \text{span}\{E_{33}, E_{22} + E_{13}\}$. Then $f_P^0 = \text{cone}\{E_{33}\}$ and $f_D^0 = \text{cone}\{E_{11}\}$. Therefore, $f_P^0 \subset (f_D^0)^c$ and $f_D^0 \subset (f_P^0)^c$, i.e., this is not a strict complementarity partition. In addition, note that (2.19) holds and moreover, if we choose $\tilde{s} = \tilde{x} = E_{22} \in (f_P^0)^c \cap (f_D^0)^c$, then $E_{22} \in \text{precl}(\mathcal{L} + (\text{face } \tilde{s})^c) \cap \text{precl}(\mathcal{L}^\perp + (\text{face } \tilde{s})^c)$. This means that $0 \neq \bar{s} = \bar{x} = E_{22}$ is both primal and dual optimal, see Theorem 5.3, below. Since $\langle \bar{s}, \bar{x} \rangle > 0$, we have obtained a finite positive duality gap.

Example 2.6 We can use $K = K^* = \mathbb{S}_+^n$ and the algorithm in [71] to generate \mathcal{A} so that we have $(f_P^0)^c \cap (f_D^0)^c \neq \{0\}$. Here the linear transformation $\mathcal{A}^* y = \sum_{i=1}^m y_i A_i$ for given $A_i \in \mathbb{S}^n, i = 1, \dots, m$. The main idea is to start with $[Q_P \ Q_G \ Q_D]$ an orthogonal matrix; and then we construct one of the m matrices representing \mathcal{A} as

$$A_1 = [Q_P \ Q_G \ Q_D] \begin{bmatrix} 0 & 0 & Y_2^T \\ 0 & \boxed{Y_1} & Y_3^T \\ Y_2 & Y_3 & Y_4 \end{bmatrix} [Q_P \ Q_G \ Q_D]^T,$$

where $Y_1 \succ 0$, Y_4 symmetric, and $Q_D Y_2 \neq 0$. The other matrices $A_i \in \mathbb{S}^n$ are chosen so that the set $\{A_1 Q_P, \dots, A_m Q_P\}$ is linearly independent. Then we get the partition in positions given by

$$\begin{bmatrix} f_P^0 & \cdot & \cdot \\ \cdot & G & \cdot \\ \cdot & \cdot & f_D^0 \end{bmatrix},$$

where G represents the gap in the partition.

All instances in the above examples have the facial block structure $\begin{bmatrix} f_D^0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & f_P^0 \end{bmatrix}$. Viz., the matrices in f_D^0 are nonzero only in the (1, 1) block, and the matrices in f_P^0 are nonzero only in the (3, 3) block. We now formalize the concept of such block structure for \mathbb{S}^n in the following definition and lemma. (These may be extended to more general cones using appropriate bases.)

Definition 2.7 The support of $x \in \mathbb{S}^n$ is $\mathcal{S}(x) := \{(ij) : x_{ij} \neq 0\}$.

Lemma 2.8 Let $K := \mathbb{S}_+^n$.

1. There exists an orthogonal matrix Q and integers $0 \leq k_D < k_P \leq n + 1$ such that

$$x \in f_D^0, (ij) \in \mathcal{S}(Q^T x Q) \implies \max\{i, j\} \leq k_D, \quad (2.20)$$

and

$$s \in f_P^0, (ij) \in \mathcal{S}(Q^T s Q) \implies \min\{i, j\} \geq k_P. \quad (2.21)$$

2. Let $n \geq 3$ and suppose the subspace \mathcal{L} is such that the complementarity partition (f_P^0, f_D^0) is not strict. Then, there exists an orthogonal matrix Q and integers $1 \leq k_D < k_P - 1 \leq n - 1$ such that (2.20) and (2.21) hold.

Proof.

We can choose $x = Q_x D_x Q_x^T \in \text{relint } f_D^0$ and $s = Q_s D_s Q_s^T \in \text{relint } f_P^0$, where Q_x, Q_s has orthonormal columns (of eigenvectors) and both D_x and D_s are diagonal positive definite. Choose Q_r so that $Q := [Q_x \ Q_r \ Q_s]$ is an orthogonal matrix. Then this Q does what we want, since $f_P^0 f_D^0 = 0$. ■

3 Facial Properties

We now collect some interesting though technical facial properties for general convex cones K . These results are particularly useful for SDP. We include the notions of nice and devious cones. Further results are given in e.g., [7, 8, 17, 51].

3.1 Faces of General Cones

Recall that a nonempty face $F \trianglelefteq K$ is *exposed* if $F = \phi^\perp \cap K$, for some $\phi \in K^*$. Note that the faces of K are closed if K is closed.

Proposition 3.1 *Let K be closed and $\emptyset \neq F \trianglelefteq K$. Then:*

1. $(F - F) \cap K = (\text{span } F) \cap K = F$.
2. $F^{cc} = F$ if and only if F is exposed.
3. $\overline{K^* + \text{span } F^c} \subseteq \overline{K^* + F^\perp}$. Moreover, if K is facially exposed, then $\overline{K^* + \text{span } F^c} = \overline{K^* + F^\perp}$.

Proof.

1. That $F - F = \text{span } F$ follows from the definition of a cone. Further, suppose $s = f_1 - f_2$ with $s \in K$ and $f_i \in F, i = 1, 2$. Then $s + f_2 = f_1 \in F$. Therefore, $s \in F$, by the definition of a face.
2. The result follows from the facts: the conjugate of $G := F^c$ is exposed by any $x \in \text{relint } G$; and, every exposed face is exposed by any point in the relative interior of its conjugate face.
3. That $K^* + \text{span } F^c \subseteq K^* + F^\perp$ is clear from the definition of the conjugate face F^c . To prove equality, suppose that K is facially exposed and that $w = (x + f) \in (K^* + F^\perp) \setminus \overline{K^* + \text{span } F^c}$, with $x \in K^*, f \in F^\perp$. Then there exists ϕ such that $\langle \phi, w \rangle < 0 \leq \langle \phi, (K^* + \text{span } F^c) \rangle$. This implies that $\phi \in K \cap (F^c)^\perp = K \cap F$, since K is facially exposed. This in turn implies $\langle \phi, w \rangle = \langle \phi, x + f \rangle \geq 0$, a contradiction. ■

Proposition 3.2 *Let $s \in \text{relint } S$ and $S \subseteq K$ be a convex set. Then:*

1. $\text{face } s = \text{face } S$,
2. $\text{cone}(K - s) = \text{cone}(K - S) = K - \text{face } s = K - \text{face } S = K + \text{span } \text{face } s = K + \text{span } \text{face } S$.

Proof.

1. That $\text{face } s \subseteq \text{face } S$ is clear. To prove the converse inclusion, suppose that $z \in S \subseteq K, z \neq s$. Since $s \in \text{relint } S$, there exists $w \in S, 0 < \theta < 1$, such that $s = \theta w + (1 - \theta)z$, i.e., $s \in (w, z)$. Since $s \in \text{face } s$, we conclude that both $w, z \in \text{face } s$.
2. That $\text{cone}(K - s) \subseteq K - \text{cone } s \subseteq K - \text{face } s \subseteq K - \text{span } \text{face } s$ is clear. The other inclusions follow from part 1 and $\text{cone}(K - s) \supseteq \text{cone}(\text{face}(s) - s) = \text{span } \text{face } s$. ■

We can combine Proposition 2.4 and Proposition 3.2, and conclude that K is polyhedral if and only if $K + \text{span } F$ is closed, for all $F \trianglelefteq K$. The following Proposition 3.3 illustrates some technical properties of faces, conjugates, and closure.

Proposition 3.3 *Let T be a convex cone and $F \trianglelefteq T$.*

1. Suppose that $\bar{s} \in \text{relint } F$ and $\bar{x} \in \text{relint } F^c$. Then

$$\bar{s} + \bar{x} \in \text{int } (T + T^*).$$

2. Suppose that $\bar{s} \in \text{relint } F$. Then

$$\begin{aligned} \overline{\text{cone}(T - \bar{s})} &= (F^c)^*; \\ \text{cone}(T - \bar{s}) &\supseteq \text{relint } ((F^c)^*). \end{aligned} \tag{3.1}$$

Proof.

1. First, note that if $\text{int } (T + T^*) = \emptyset$, then we have

$$0 \neq (T + T^*)^\perp \subseteq T^{**} \cap T^* = \bar{T} \cap T^* \subseteq T + T^*,$$

a contradiction, i.e., this shows that $\text{int } (T + T^*) \neq \emptyset$.

Now suppose that $\bar{s} + \bar{x} \notin \text{int } (T + T^*)$. Then we can find a supporting hyperplane ϕ^\perp so that $\bar{s} + \bar{x} \in (T + T^*) \cap \phi^\perp \triangleleft T + T^*$ and $0 \neq \phi \in (T + T^*)^* = \bar{T} \cap T^*$. Therefore, we conclude $\langle \phi, \bar{s} + \bar{x} \rangle = 0$ implies that both $\langle \phi, \bar{s} \rangle = 0$ and $\langle \phi, \bar{x} \rangle = 0$. This means that $\phi \in T^* \cap \bar{s}^\perp = F^c$ and $\phi \in \bar{T} \cap \bar{x}^\perp = (F^c)^c$, giving $0 \neq \phi \in (F^c) \cap (F^c)^c = \{0\}$, which is a contradiction.

2. The first result follows from: $\overline{\text{cone}(T - \bar{s})} = (T - \bar{s})^{**} = (T^* \cap \bar{s}^\perp)^*$.

For the second result, we use the first result and Theorem 6.3 of Rockafellar [63] to deduce

$$\text{relint } (\text{cone}(T - \bar{s})) = \text{relint } \left(\overline{\text{cone}(T - \bar{s})} \right) = \text{relint } [(F^c)^*]$$

which implies the desired conclusion. ■

3.2 Faces for Primal-Dual Pair (P) and (D)

We now present facial properties specific to the primal-dual pair (2.2) and (2.3). In particular, this includes relationships between the minimal faces f_P, f_D and the minimal faces for the homogeneous problems, f_P^0, f_D^0 .

Proposition 3.4 *Suppose that both (2.2) and (2.3) are feasible, i.e., equivalently $\tilde{s} \in K + \mathcal{L}^\perp$ and $\tilde{x} \in K^* + \mathcal{L}$. Let $\hat{s} \in \mathcal{F}_P(\tilde{s})$ and $\hat{x} \in \mathcal{F}_D(\tilde{x})$. Then the following hold.*

1.

$$f_P^0 \subseteq \text{face}(\hat{s} + f_P^0) \subseteq f_P(\tilde{s}), \quad f_D^0 \subseteq \text{face}(\hat{x} + f_D^0) \subseteq f_D(\tilde{x}). \tag{3.2}$$

2.

$$\tilde{s} \in (f_D^0)^c + \mathcal{L}^\perp \Leftrightarrow f_P(\tilde{s}) \subseteq (f_D^0)^c; \quad \tilde{x} \in (f_P^0)^c + \mathcal{L} \Leftrightarrow f_D(\tilde{x}) \subseteq (f_P^0)^c.$$

Proof.

Since both problems are feasible, we can assume, without loss of generality, that $\hat{s} = \tilde{s} \in K, \hat{x} = \tilde{x} \in K^*$.

1. Since cone \tilde{s} and f_P^0 are convex cones containing the origin, cone $\tilde{s} + f_P^0 = \text{conv}(\text{cone } \tilde{s} \cup f_P^0)$; see e.g., [63, Theorem 3.8]. Hence,

$$f_P^0 \subseteq \text{conv}(\tilde{s} \cup f_P^0) \subseteq \text{conv}(\text{cone } \tilde{s} \cup f_P^0) = \text{cone } \tilde{s} + f_P^0 = \text{cone } (\tilde{s} + f_P^0) \subseteq \text{face } (\tilde{s} + f_P^0).$$

This proves the first inclusion. It is clear that $\tilde{s} + (\mathcal{L}^\perp \cap K) \subseteq (\tilde{s} + \mathcal{L}^\perp) \cap K$. This yields the second inclusion. The final two inclusions follow similarly.

2. Suppose that $\tilde{s} \in (f_D^0)^c + \mathcal{L}^\perp$ and $\tilde{s} + r \in K$ with $r \in \mathcal{L}^\perp$. Then, for all $\ell \in \mathcal{L} \cap K^* \subseteq f_D^0$, we have $\langle \tilde{s} + r, \ell \rangle = \langle \tilde{s}, \ell \rangle = 0$, since $\tilde{s} \in (f_D^0)^c + \mathcal{L}^\perp$. This implies that $f_P(\tilde{s}) = \text{face } ((\tilde{s} + \mathcal{L}^\perp) \cap K)$ is orthogonal to $f_D^0 = \text{face } (\mathcal{L} \cap K^*)$, i.e., the first implication holds. (This also follows from part 1, using $f_P(\tilde{s}) \subseteq [f_D(\tilde{x})]^c$.)

For the converse implication, since (2.2) is feasible, we have $\tilde{s} \in f_P(\tilde{s})$. So if $f_P(\tilde{s}) \subseteq (f_D^0)^c$, then $\tilde{s} \in (f_D^0)^c$.

The second equivalence follows similarly. ■

Additional relationships between the faces follow. First, we need a lemma that is of interest in its own right.

Lemma 3.5 *Let $\tilde{s} \in f_P^0$, and suppose that $s = \tilde{s} + \ell$ is feasible for (2.2) with $\ell \in \mathcal{L}^\perp$. Then $\ell \in \text{span } f_P^0$.*

Proof.

Let $v \in \text{relint } (\mathcal{L}^\perp \cap K)$. Then $v \in \text{relint } f_P^0$, and since $\tilde{s} \in f_P^0$, we have $v - \epsilon \tilde{s} \in f_P^0$ for some $\epsilon > 0$. Now if ℓ is such that $s = \tilde{s} + \ell$ is feasible for (2.2), then $\tilde{s} + \ell \in K$, and

$$\frac{1}{\epsilon}v + \ell = \frac{1}{\epsilon}(v - \epsilon \tilde{s}) + (\tilde{s} + \ell) \in f_P^0 + K = K.$$

For convenience, define $\alpha := 1/\epsilon$. Since $\alpha v \in \mathcal{L}^\perp$ and $\ell \in \mathcal{L}^\perp$, we in fact have

$$\alpha v + \ell \in K \cap \mathcal{L}^\perp \subseteq f_P^0, \tag{3.3}$$

which implies $\ell \in f_P^0 - f_P^0$. ■

Proposition 3.6 1. $\tilde{s} \in f_P^0 \cup \mathcal{L}^\perp \implies f_P(\tilde{s}) = f_P^0$ and $\tilde{x} \in f_D^0 \cup \mathcal{L} \implies f_D(\tilde{x}) = f_D^0$.

2. Let $f_D^0 \triangleleft K^*$. Then there exists $0 \neq \phi \in K \cap \mathcal{L}^\perp$.

Proof.

1. We begin by proving the first statement. If $\tilde{s} \in \mathcal{L}^\perp$, then $\tilde{s} + \mathcal{L}^\perp = \mathcal{L}^\perp$, so the desired result holds. If instead $\tilde{s} \in f_P^0$, then it follows from Lemma 3.5 that $\ell \in \text{span } f_P^0$ for all feasible points of the form $s = \tilde{s} + \ell$. Hence all feasible s lie in the set $\text{span } (f_P^0) \cap K$, which by Proposition 3.1, part 1, equals f_P^0 . So $f_P(\tilde{s}) \subseteq f_P^0$; but, the reverse inclusion holds by Proposition 3.4, part 1.

The second statement for f_D^0 is proven in a similar way.

2. Existence is by the theorem of the alternative for the Slater CQ; see (2.15) and the related Proposition 2.3. ■

We conclude this subsection with a new result indicating that the failure of strict complementarity in (f_P^0, f_D^0) can be related to the lack of closure in the sum of the cone and the subspace.

Proposition 3.7 *Let K be closed. If there exists a nonzero $x \in -(K \cap K^*)$ such that*

$$x \in (K \cap \mathcal{L}^\perp)^\perp \cap (K^* \cap \mathcal{L})^\perp, \quad (3.4)$$

then $x \in \text{precl}(K + \mathcal{L}^\perp) \cap \text{precl}(K^ + \mathcal{L})$. Hence, neither $K + \mathcal{L}^\perp$ nor $K^* + \mathcal{L}$ is closed.*

Proof.

To obtain a contradiction, suppose that (3.4) holds for a nonzero $x \in -(K \cap K^*)$, but $x \in K + \mathcal{L}^\perp$. Then there exists $w \in K$ such that $x - w \in \mathcal{L}^\perp$. Moreover, $x - w \in -K - K = -K$, so $x - w \in -(K \cap \mathcal{L}^\perp)$. It follows from (3.4) that $\langle x, x - w \rangle = 0$. However, $\langle x, x - w \rangle = \langle x, x \rangle + \langle -x, w \rangle > 0$, where we have used the fact that $x \in -K^*$. Hence, $x \notin K + \mathcal{L}^\perp$. A similar argument shows that $x \notin K^* + \mathcal{L}$.

Since K and \mathcal{L}^\perp are closed convex cones, we have $(K \cap \mathcal{L}^\perp)^* = \overline{K^* + (\mathcal{L}^\perp)^*} = \overline{K^* + \mathcal{L}}$. It follows from (3.4) that $x \in \overline{K^* + \mathcal{L}}$. Similarly, $x \in \overline{K + \mathcal{L}^\perp}$. This completes the proof. ■

The reader might naturally ask: when does such an x satisfying the conditions of the last proposition exist? As an answer, consider the case $K = K^*$ and a linear subspace \mathcal{L} such that (f_P^0, f_D^0) is a proper partition but is not strictly complementary. Let $G := (f_P^0)^c \cap (f_D^0)^c$. Then every $x \in -G \setminus \{0\}$ satisfies the assumption of the above proposition.

3.3 Nice Cones, Devious Cones, and SDP

Definition 3.8 *A face $F \triangleleft K$ is called nice if $K^* + F^\perp$ is closed. A closed convex cone K is called a nice cone or a facially dual-complete cone, FDC, if*

$$K^* + F^\perp \text{ is closed for all } F \triangleleft K. \quad (3.5)$$

The condition in (3.5) was used in [13] to allow for extended Lagrange multipliers in f_P^* to be split into a sum using K^* and f_P^\perp . This allowed for restricted Lagrange multiplier results with the multiplier in K^* . The condition (3.5) was also used in [52] where the term *nice cone* was introduced. In addition, it was shown by Pataki (forthcoming paper) that a FDC cone must be facially exposed.

Moreover, the FDC property has an implication for Proposition 3.1, part 3. We now see that this holds for SDP.

Lemma 3.9 ([73],[58]) *Suppose that F is a proper face of \mathbb{S}_+^n , i.e., $\{0\} \neq F \triangleleft \mathbb{S}_+^n$. Then:*

$$F^* = \mathbb{S}_+^n + F^\perp = \overline{\mathbb{S}_+^n + \text{span } F^c},$$

$$\mathbb{S}_+^n + \text{span } F^c \text{ is not closed.}$$

■

From Lemma 3.9, we see that \mathbb{S}_+^n is a nice cone. In fact, as pointed out in [52], many other classes of cones are nice cones, e.g., polyhedral and p-cones. However, the lack of closure property in Lemma 3.9 is *not* a nice property. In fact, from Proposition 3.2, part 2, this corresponds to the lack of closure for radial cones, see [66] which can result in duality problems. Therefore, we add the following.

Definition 3.10 *A face $F \trianglelefteq K$ is called devious if the set $K + \text{span } F$ is not closed. A cone K is called devious if*

$$\text{the set } K + \text{span } F \text{ is not closed for all } \{0\} \neq F \triangleleft K.$$

By Lemma 3.9, \mathbb{S}_+^n is a nice but devious cone. On the other hand, polyhedral cones are nice and not devious, since sums of polyhedral cones and subspaces are closed, e.g., [63, Chapter 9].

The facial structure of \mathbb{S}_+^n is well known, e.g., [58, 73]. Each face $F \trianglelefteq \mathbb{S}_+^n$ is characterized by a unique subspace $S \subseteq \mathbb{R}^n$:

$$F = \{x \in \mathbb{S}_+^n : \mathcal{N}(x) \supseteq S\}; \quad \text{relint } F = \{x \in \mathbb{S}_+^n : \mathcal{N}(x) = S\}.$$

The conjugate face satisfies

$$F^c = \{x \in \mathbb{S}_+^n : \mathcal{N}(x) \supseteq S^\perp\}; \quad \text{relint } F^c = \{x \in \mathbb{S}_+^n : \mathcal{N}(x) = S^\perp\}.$$

The description of $\text{span } F$ for $F \trianglelefteq \mathbb{S}_+^n$ is now clear.

Another useful property of SDPs (and the Löwner partial order) is given by the following lemma. This lemma played a critical role in the explicit description of a dual SDP problem for which strong duality holds.

Lemma 3.11 ([57]) *Let $\tilde{K} \subseteq \mathbb{S}_+^n$ be a closed convex cone. Then*

$$\left[\left(\text{face } \tilde{K} \right)^c \right]^\perp = \left\{ W + W^T : W \in \mathbb{R}^{n \times n}, \begin{bmatrix} I & W^T \\ W & U \end{bmatrix} \succeq 0, \text{ for some } U \in \tilde{K} \right\}.$$

■

Properties 3.12 *The following three properties of the cone \mathbb{S}_+^n are needed for the strong duality approach in Ramana [57]. The first two also make the Borwein-Wolkowicz approach in [15] behave particularly well:*

1. K is facially exposed.
2. K is FDC.
3. Lemma 3.11.

■

Suppose that the cone K describing the problem (P) is *SDP-representable*. (That is, there exists d and a linear subspace $V \subset \mathbb{S}^d$ such that $V \cap \mathbb{S}_{++}^d \neq \emptyset$ and K is isomorphic to $(V \cap \mathbb{S}_+^d)$.) Then by [20, Cor. 1, Prop. 4], K is facially exposed and FDC, since \mathbb{S}_+^d is. Moreover, by [20, Prop. 3], every proper face of K is a proper face of \mathbb{S}_+^d intersected with the subspace V . Hence, assuming that a suitable representation of K is given, an analogue of Lemma 3.11 is also available in this case. Therefore, SDP-representable cones (which strictly include homogeneous cones, due to a result of Chua [19] and Faybusovich [24]) satisfy all three of the above-mentioned Properties 3.12. For related recent results on homogeneous cones and strong duality, see Pólik and Terlaky [53].

4 Duality and Minimal Representations

In this section, we see that *minimal representations* of the problem guarantee strong duality and stability results, i.e., combining the minimal cone and the minimal subspace together reduces both the dimension of the problem and the number of constraints, and also guarantees Slater's constraint qualification. We first use the minimal subspaces and extend the known strong duality results without any constraint qualification, see e.g., [13, 14, 15, 72]. Equivalent strong duality results based on an extended Lagrangian are given in [56, 57]. (See [58, 73] for comparison and summaries of the two types of duality results.) By strong duality for (2.2), we mean that there is a *zero duality gap*, $v_P = v_D$, and the dual optimal value v_D in (2.3) is attained.

4.1 Strong Duality and Constraint Qualifications

We now present strong duality results that hold with and without CQs. We also present: a weakest constraint qualification (WCQ), i.e., a CQ at a given feasible point $\bar{y} \in \mathcal{F}_P^y(c)$ that is independent of b ; and a universal constraint qualification, (UCQ), i.e., a CQ that is independent of both b and c . Following is the classical, well-known, strong duality result for (2.2) under the standard Slater CQ.

Theorem 4.1 (e.g., [46, 65]) *Suppose that Slater's CQ (strict feasibility) holds for (2.2). Then strong duality holds for (2.2), i.e., $v_P = v_D$ and the dual value v_D in (2.3) is attained. Equivalently, there exists $\bar{x} \in K^*$ such that*

$$\langle b, y \rangle + \langle c - \mathcal{A}^* y, \bar{x} \rangle \geq v_P, \quad \forall y \in \mathbb{R}^m.$$

Moreover, if v_P is attained at $\bar{y} \in \mathcal{F}_P^y$, then $\langle c - \mathcal{A}^* \bar{y}, \bar{x} \rangle = 0$ (complementary slackness holds). ■

A nice compact formulation follows.

Corollary 4.2 *Suppose that Slater's CQ (strict feasibility) holds for (2.2) and $\bar{y} \in \mathcal{F}_P^y$. Then, \bar{y} is optimal for (2.2) if and only if*

$$b \in \mathcal{A}[(K - \bar{s})^*], \tag{4.1}$$

where $\bar{s} = c - \mathcal{A}^* \bar{y}$.

Proof.

The result follows from the observation that $(\text{face } \bar{s})^c = K^* \cap \bar{s}^\perp = (K - \bar{s})^*$, i.e., (4.1) is equivalent to dual feasibility and complementary slackness. ■

Strong duality can fail if Slater's CQ does not hold. In [14, 13, 15], an equivalent *regularized* primal problem that is based on the minimal face,

$$v_{RP} := \sup\{\langle b, y \rangle : \mathcal{A}^* y \preceq_{f_P} c\} \tag{4.2}$$

is considered. Its Lagrangian dual is given by

$$v_{DRP} := \inf\{\langle c, x \rangle : \mathcal{A}x = b, x \succeq_{f_P^*} 0\}. \tag{4.3}$$

Theorem 4.3 ([13]) *Strong duality holds for the pair (4.2) and (4.3), or equivalently, for the pair (2.2) and (4.3); i.e., $v_P = v_{RP} = v_{DRP}$ and the dual optimal value v_{DRP} is attained. Equivalently, there exists $x^* \in (f_P)^*$ such that*

$$\langle b, y \rangle + \langle c - \mathcal{A}^* y, x^* \rangle \geq v_P, \quad \forall y \in (\mathcal{A}^*)^{-1}(f_P - f_P).$$

Moreover, if v_P is attained at $\bar{y} \in \mathcal{F}_P^y$, then $\langle c - \mathcal{A}^* \bar{y}, x^* \rangle = 0$ (complementary slackness holds). ■

Corollary 4.4 *Let $\bar{y} \in \mathcal{F}_P^y$. Then \bar{y} is optimal for (2.2) if and only if*

$$b \in \mathcal{A}[(f_P - \bar{s})^*],$$

where $\bar{s} = c - \mathcal{A}^* \bar{y}$.

Proof.

As above, in the proof of Corollary 4.2, the result follows from the observation that $f_P^* \cap \bar{s}^\perp = (f_P - \bar{s})^*$. ■

The next result uses the *minimal subspace representation* of \mathcal{L}^\perp , introduced in (2.17), $\mathcal{L}_{PM}^\perp = \mathcal{L}^\perp \cap (f_P - f_P)$.

Corollary 4.5 *Let \tilde{y}, \tilde{s} , and \tilde{x} satisfy (2.7) with $\tilde{s} \in f_P - f_P$ and let*

$$K^* + (f_P)^\perp = (f_P)^*. \quad (4.4)$$

Consider the following pair of dual programs.

$$v_{RP_M} = \langle c, \tilde{x} \rangle - \inf_s \{ \langle s, \tilde{x} \rangle : s \in (\tilde{s} + \mathcal{L}_{PM}^\perp) \cap K \}, \quad (4.5)$$

$$v_{DRP_M} = \langle b, \tilde{y} \rangle + \inf_x \{ \langle \tilde{s}, x \rangle : x \in (\tilde{x} + \mathcal{L}_{PM}) \cap K^* \}. \quad (4.6)$$

Then, $v_{RP_M} = v_{RP} = v_P = v_{DRP_M} = v_{DRP}$, and strong duality holds for (4.5) and (4.6), or equivalently, for the pair (2.2) and (4.6).

Proof.

That $v_P = v_{RP_M} = v_{RP}$ follows from the definition of the minimal subspace representation in (2.17):

$$\begin{aligned} \mathcal{F}_P^s(c) &= \mathcal{F}_P^s(\tilde{s}) \\ &= (\tilde{s} + \mathcal{L}^\perp) \cap f_P, \quad \text{by definition of } f_P, \\ &= (\tilde{s} + \mathcal{L}_{PM}^\perp) \cap K, \quad \text{since } \tilde{s} \in f_P - f_P. \end{aligned}$$

For the regularized dual, we see that

$$\begin{aligned} v_{DRP} &= \inf_x \left\{ \langle c, x \rangle : \mathcal{A}x = b, x \succeq_{f_P^*} 0 \right\} \\ &= \langle \tilde{y}, b \rangle + \inf_x \left\{ \langle \tilde{s}, x \rangle : \mathcal{A}x = b, x = x_k + x_f, x_k \in K^*, x_f \in f_P^\perp \right\}, \quad \text{by (4.4)} \\ &= \langle \tilde{y}, b \rangle + \inf_x \left\{ \langle \tilde{s}, x \rangle : x = x_k + x_f = \tilde{x} + x_l, x_k \in K^*, x_f \in f_P^\perp, x_l \in \mathcal{L} \right\} \\ &= \langle \tilde{y}, b \rangle + \inf_x \left\{ \langle \tilde{s}, x_k \rangle : x_k \in (\tilde{x} + \mathcal{L} + f_P^\perp) \cap K^* \right\} = v_{DRP_M}. \end{aligned}$$

■

Remark 4.6 *The condition in (4.4) is equivalent to $K^* + (f_P)^\perp$ being closed, and is clearly true for every face of K , if K is a FDC cone.*

Remark 4.7 *Using the minimal subspace representations of \mathcal{L} in (2.3), i.e., replacing \mathcal{L} in (2.3) by \mathcal{L}_{DM} in (2.17), we may obtain a result similar to Corollary 4.5.*

Note that if the Slater CQ holds, then the minimal sets (face and subspace) satisfy $f_P = K$ and (2.17). We now see that if at least one of these conditions holds, then strong duality holds.

Corollary 4.8 *Suppose that $\text{int } K = \emptyset$ but the generalized Slater CQ (relative strict feasibility) holds for (2.2), i.e.,*

$$\hat{s} := c - \mathcal{A}^* \hat{y} \in \text{relint } K, \text{ for some } \hat{y} \in \mathcal{W}. \quad (\text{Generalized Slater CQ}) \quad (4.7)$$

(Equivalently, suppose that the minimal face satisfies $f_P = K$.) Then strong duality holds for (2.2).

Proof.

The proof follows immediately from Theorem 4.3 after noting that $K = f_P$. ■

The following corollary illustrates strong duality for a variation of the generalized Slater constraint qualification, i.e., for the case that the minimal subspace satisfies (2.17).

Corollary 4.9 *Let $\tilde{s} \in f_P - f_P$ and K be FDC. Suppose that*

$$\mathcal{L}^\perp \cap (K - K) \subseteq f_P - f_P. \quad (\text{Subspace CQ}) \quad (4.8)$$

(Equivalently, suppose that $\mathcal{L}_{P_M}^\perp = \mathcal{L}^\perp \cap (K - K)$.) Then strong duality holds for (2.2).

Proof.

Follows directly from Corollary 4.5. ■

We now summarize the results in the special case that K is FDC (a nice cone). The first item presents a regularized problem that satisfies Slater's CQ. This is the approach used in [18]. Note that early results on weakest constraint qualifications for general nonlinear problems are given in e.g., [31].

Theorem 4.10 *Let \tilde{s}, \tilde{x} satisfy linear feasibility (2.7) with $\tilde{s} \in f_P - f_P$ and let K be FDC. Then we have the following conclusions.*

1. *The primal optimal values are all equal, $v_P = v_{RP} = v_{RP_M}$. Moreover, strong duality holds for the primal, where the primal is chosen from the set*

$$\{(2.9), (4.2), (4.5)\} \quad (\text{set of primal programs})$$

and the dual is chosen from the set

$$\{(4.3), (4.6)\} \quad (\text{set of dual programs})$$

i.e., the optimal values are all equal and the dual optimal value is attained.

2. Furthermore, let $\mathcal{T} : \mathbb{R}^t \rightarrow \mathcal{V}$ be a one-one linear transformation with $\mathcal{R}(\mathcal{T}) = f_P - f_P$. Then Slater's CQ holds for the regularized problem

$$v_{RP_M} = \langle c, \tilde{x} \rangle - \inf_{v \in \mathbb{R}^t} \{ \langle v, \mathcal{T}^* \tilde{x} \rangle : v \in (\mathcal{T}^\dagger \bar{s} + \mathcal{T}^\dagger(\mathcal{L}_{P_M}^\perp)) \cap \mathcal{T}^\dagger(f_P) \}. \quad (4.9)$$

3. The following are CQs for (2.2):

- (a) $f_P = K$ (equivalently generalized Slater CQ (4.7));
(b) $\mathcal{L}^\perp \cap (K - K) \subseteq f_P - f_P$ (equivalently $\mathcal{L}_{P_M}^\perp = \mathcal{L}^\perp \cap (K - K)$);

4. Let $\bar{y} \in \mathcal{F}_P^y(c)$ and $\bar{s} = c - \mathcal{A}^* \bar{y}$. Then,

$$\mathcal{D}_{\bar{P}}^{\leq}(\bar{y})^* = -\mathcal{A}((K - \bar{s})^*) \quad \text{is a WCQ for (2.2) at } (\bar{y}, \bar{s}). \quad (4.10)$$

Equivalently,

$$\mathcal{A}[(f_P - \bar{s})^*] = \mathcal{A}((K - \bar{s})^*) \quad \text{is a WCQ for (2.2) at } (\bar{y}, \bar{s}). \quad (4.11)$$

Proof.

1. These results summarize Theorem 4.3 and Corollary 4.5.
2. From the definitions, we know that there exists $\ell \in \mathcal{L}_{P_M}^\perp$ with $\hat{s} = \bar{s} + \ell \in \text{relint } f_P$. Therefore, $v := \mathcal{T}^\dagger \hat{s} = \mathcal{T}^\dagger(\bar{s} + \ell) \in \text{int } \mathcal{T}^\dagger(f_P)$.
3. The results follow from Corollaries 4.8, 4.9, respectively.
4. The so-called Rockafellar-Pshenichnyi condition, e.g., [55], [31, Thm 1], states that \bar{y} is optimal if and only if $b \in -\mathcal{D}_{\bar{P}}^{\leq}(\bar{y})^*$. From Theorem 4.3, \bar{y} is optimal if and only if $\mathcal{A} \bar{x} = b$, $\langle \bar{s}, \bar{x} \rangle = 0$, for some $\bar{x} \in f_P^*$; equivalently, if and only if $b \in \mathcal{A}((f_P - \bar{s})^*)$. The result follows from the fact that strong duality holds at an optimal \bar{y} if and only if $\mathcal{A} \bar{x} = b$, $\langle \bar{s}, \bar{x} \rangle = 0$, for some $\bar{x} \in K^*$; equivalently, $\mathcal{A} \bar{x} = b$, for some $\bar{x} \in (K - \bar{s})^*$.

■

Remark 4.11 The WCQ in (4.10) follows the approach in e.g., [31, 22, 76, 43, 44]. Moreover, since for any set S , $\mathcal{A}(S)$ is closed if and only if $S + \mathcal{L}$ is closed (e.g., [34, 10]), we conclude that a necessary condition for the WCQ to hold at a feasible $\bar{s} \in \mathcal{F}_P^s$ is that

$$(K - \bar{s})^* + \mathcal{L} \quad \text{is closed.} \quad (4.12)$$

(For a recent detailed study of when the linear image of a closed convex set is closed, see e.g., [52]. For related perturbation results, see [12].)

4.1.1 Universal Constraint Qualifications

A universal CQ, denoted UCQ, is a CQ that holds independent of the data b, c , i.e., as in LP, strong duality holds for arbitrary perturbations of the data b, c as long as feasibility is not lost.

Theorem 4.12 *Suppose that K is FDC, and $\tilde{s} \in K, \tilde{x} \in K^*$ in the primal-dual subspace representation in (2.9) and (2.10). Then*

$$\mathcal{L}^\perp \subseteq f_P^0 - f_P^0$$

is a UCQ, i.e., a universal CQ for (2.2).

Proof.

The result follows from Corollary 4.9 and the fact that $\tilde{s} \in K, \tilde{x} \in K^*$ implies $f_P^0 \subseteq f_P$ and $f_D^0 \subseteq f_D$, see Proposition 3.4. ■

Corollary 4.13 *Suppose that $K = \mathbb{S}_+^n$, both v_P and v_D are finite, and $n \leq 2$. Then strong duality holds for at least one of (2.2) or (2.3).*

Proof.

We have both $\mathcal{F}_P^s \neq \emptyset$ and $\mathcal{F}_D^x \neq \emptyset$. By going through the possible cases, we see that one of the CQs $\mathcal{L}^\perp \subseteq f_P^0 - f_P^0$ or $\mathcal{L} \subseteq f_D^0 - f_D^0$ must hold. ■

5 Failure of Strong Duality and Strict Complementarity

As discussed above, the absence of a CQ for (2.2) can result in the failure of strong duality, i.e., we have theoretical difficulties. In addition, it has been shown that near loss of Slater's CQ is closely correlated with an increase in the expected number of iterations in interior-point methods both in theory [59, 61] and empirically, [25, 26]. Therefore, a regularization step should be an essential preprocessor for SDP solvers.

It is also known that the lack of strict complementarity for SDP may result in theoretical difficulties. For example, superlinear and quadratic convergence results for interior-point methods depend on the strict complementarity assumption, e.g., [54, 37, 3, 47, 40]. This is also the case for convergence of the central path to the analytic center of the optimal face, [32]. In addition, it is shown empirically in [71] that the loss of strict complementarity is closely correlated with the expected number of iterations in interior-point methods. However, one can generate problems where strict complementarity fails independent of whether or not Slater's CQ holds for the primal and/or the dual, [71]. Therefore, we see a connection between the theoretical difficulty from an absence of Slater's CQ and numerical algorithms, and a similar connection for the absence of strict complementarity. We see below that duality and strict complementarity of the homogeneous problem have a surprising theoretical connection as well.

Strong duality for (2.2) means a zero duality gap, $v_P = v_D$ and dual attainment, $v_D = \langle c, x^* \rangle$, for some $x^* \in \mathcal{F}_D^x$. The CQs (resp. UCQs), introduced above in Section 4, guarantee that strong duality holds independent of the data b (resp. b and c). Under our assumption that v_P is finite valued, there are three cases of failure to consider: (i) a zero duality gap but no dual attainment; (ii) an infinite duality gap (dual infeasibility); (iii) a finite positive duality gap.

5.1 Finite Positive Duality Gaps

5.1.1 Positive Gaps and Closure Conditions

We present characterizations for a finite positive duality gap under attainment assumptions in Theorem 5.3 and Corollary 5.4. We first give sufficient conditions for a positive duality gap using well known optimality conditions based on feasible directions.

Proposition 5.1 *Let $\tilde{s} \in \mathcal{F}_P^s$, $\tilde{x} \in \mathcal{F}_D^x$, and $\langle \tilde{s}, \tilde{x} \rangle > 0$. Suppose that $\tilde{s} \in \mathcal{D}_D^{\leq}(\tilde{x})^*$ and $\tilde{x} \in \mathcal{D}_P^{\leq}(\tilde{s})^*$. Then \tilde{s} is optimal for (2.2), \tilde{x} is optimal for (2.3), and $-\infty < v_P < v_D < \infty$.*

Proof.

The optimality of \tilde{s} and \tilde{x} follows immediately from the definition of the cones of feasible directions and the Rockafellar-Pshenichnyi condition, see e.g., the proof of Theorem 4.10. The finite positive duality gap follows from the hypotheses that both (2.2) and (2.3) are feasible, and that $\langle \tilde{s}, \tilde{x} \rangle > 0$. ■

To connect the conditions using the cones of feasible directions with conditions using closure we need the following.

Lemma 5.2 *Suppose that \tilde{y} is feasible for (2.2), with corresponding slack \tilde{s} , and that \tilde{x} is feasible for (2.3). Then*

$$-\mathcal{D}_P^{\leq}(\tilde{y})^* = \mathcal{A}((f_P - \tilde{s})^*) = \overline{\mathcal{A}((K - \tilde{s})^*)}, \quad \mathcal{D}_D^{\leq}(\tilde{x})^* = \mathcal{L}^\perp + (f_D - \tilde{x})^* = \overline{\mathcal{L}^\perp + (K^* - \tilde{x})^*}.$$

Proof.

First note that $b \in \mathcal{A}((K - \tilde{s})^*)$ implies that \tilde{s} is optimal for (2.2). Therefore, $b \in -\mathcal{D}_P^{\leq}(\tilde{y})^* = \mathcal{A}((f_P - \tilde{s})^*)$, by the Rockafellar-Pshenichnyi condition and Corollary 4.4, i.e., we have shown the first containment $\overline{\mathcal{A}((K - \tilde{s})^*)} \subseteq -\mathcal{D}_P^{\leq}(\tilde{y})^* = \mathcal{A}((f_P - \tilde{s})^*)$.

To prove the converse containment, we use Proposition 3.3, part 2, and show that $(\mathcal{A}((K - \tilde{s})^*))^* \subseteq -\mathcal{D}_P^{\leq}(\tilde{y})$. Let $F = \text{face } \tilde{s}$ and let $d \in (\mathcal{A}((K - \tilde{s})^*))^* = (\mathcal{A}(F^c))^*$. Equivalently, $\mathcal{A}^* d \in (F^c)^*$. Then Proposition 3.3, part 2, implies that $-\mathcal{A}^* d$ is in the tangent cone to \mathcal{F}_P^s at \tilde{s} ; equivalently, $-d$ is in the tangent cone to \mathcal{F}_P^y at \tilde{y} .

The second containment follows similarly. ■

A well-known characterization for a zero duality gap can be given using the perturbation function. For example, define

$$v_P(\epsilon) := \sup_y \{ \langle b, y \rangle : \mathcal{A}^* y \preceq_K c + \epsilon \}, \quad \text{where } \epsilon \in \mathcal{V}.$$

The connection with the dual functional $\phi(x) := \sup_y \langle b, y \rangle + \langle x, c - \mathcal{A}^* y \rangle$ is given in e.g., [46]. Then the geometry shows that the closure of the epigraph of v_P characterizes a zero duality gap. We now use preclosure and the optimal solution sets $\mathcal{O}_P^s, \mathcal{O}_D^x$, to present a characterization for a finite positive duality gap in the case of attainment of the primal and dual optimal values.

Theorem 5.3 *Suppose that K is closed, \tilde{y} is feasible for (2.2), with corresponding slack \tilde{s} , and that \tilde{x} is feasible for (2.3). Then*

$$\tilde{s} \in \mathcal{O}_P^s, \tilde{x} \in \mathcal{O}_D^x, \quad \langle \tilde{s}, \tilde{x} \rangle > 0,$$

if and only if

$$\tilde{x} \in \text{precl}(\mathcal{L} + (K - \tilde{s})^*), \quad c \in \text{precl}(\mathcal{L}^\perp + (K^* - \tilde{x})^*).$$

Proof.

We have: $\tilde{s} \in \mathcal{O}_P^s$ if and only if $-b \in \mathcal{D}_P^{\leq}(\tilde{y})^*$, and $\tilde{x} \in \mathcal{O}_D^x$ if and only if $c \in \mathcal{D}_D^{\leq}(\tilde{x})^*$. However, $\langle \tilde{s}, \tilde{x} \rangle > 0$ implies that strong duality fails for both the primal and the dual. Therefore, $b \notin \mathcal{A}((K - \tilde{s})^*)$ and $c \notin \mathcal{L}^\perp + (K^* - \tilde{x})^*$. The result now follows from Lemma 5.2 and the fact that $\mathcal{A}\tilde{x} = b \in \text{precl}(\mathcal{A}(K - \tilde{s})^*)$ if and only if $\tilde{x} \in \text{precl}(\mathcal{L} + (K - \tilde{s})^*)$. (See e.g., [34, 10].) ■

It is difficult to apply the closure condition in Theorem 5.3 directly. We now see that we can expand the expression under the closure using the recession cones f_P^0 and f_D^0 .

Corollary 5.4 *Suppose that K is closed and that K and K^* are facially exposed. Let $\tilde{y} = 0$ be feasible for (2.2), with corresponding slack \tilde{s} , and \tilde{x} be feasible for (2.3). Let:*

$$F_s := \text{face } \tilde{s}; F_x := \text{face } \tilde{x}; \quad T_D := F_s^c + ((f_D^0)^c)^\perp; T_P := F_x^c + ((f_P^0)^c)^\perp. \quad (5.1)$$

Then:

1.

$$\tilde{s} \in \mathcal{O}_P^s, \tilde{x} \in \mathcal{O}_D^x, \quad \langle \tilde{s}, \tilde{x} \rangle > 0, \quad (5.2)$$

if and only if

$$\tilde{x} \notin \mathcal{L} + F_s^c, \quad \tilde{s} \notin \mathcal{L}^\perp + F_x^c, \quad (5.3)$$

and

$$\tilde{x} \in \overline{F_s^c + ((\text{face}(F_s^c \cap \mathcal{L}))^c)^\perp + \mathcal{L}}, \quad \tilde{s} \in \overline{F_x^c + ((\text{face}(F_x^c \cap \mathcal{L}^\perp))^c)^\perp + \mathcal{L}^\perp}.$$

2. Moreover, if $F_s \subseteq (f_D^0)^c$ and $F_x \subseteq (f_P^0)^c$, then (5.2) holds if and only if

$$\tilde{x} \in \overline{T_D + \mathcal{L}} \setminus (\mathcal{L} + F_s^c), \quad \tilde{s} \in \overline{T_P + \mathcal{L}^\perp} \setminus (\mathcal{L}^\perp + F_x^c). \quad (5.4)$$

Proof.

We have $F_s \trianglelefteq K$ and $F_s^c \trianglelefteq K^*$. The result follows directly from the characterization in Theorem 5.3 if we use the following equivalences.

$$\begin{aligned} \overline{(K - \tilde{s})^* + \mathcal{L}} &= \overline{F_s^c + \mathcal{L}} \\ &= \overline{\text{cone}(F_s^c - (F_s^c \cap \mathcal{L})) + \mathcal{L}} \\ &= \overline{F_s^c + \text{span face}(F_s^c \cap \mathcal{L}) + \mathcal{L}}, \text{ by Proposition 3.2, Part 2, with } S = (F_s^c \cap \mathcal{L}) \\ &= \overline{F_s^c + ((\text{face}(F_s^c \cap \mathcal{L}))^c)^\perp + \mathcal{L}}, \text{ by Proposition 3.1, Part 3.} \end{aligned}$$

Moreover, if $F_s \subseteq (f_D^0)^c$, then $f_D^0 \subseteq \text{face}(F_s^c \cap \mathcal{L})$. We can then apply Proposition 3.1, Part 3 again. The equivalences for F_x follow similarly. ■

5.1.2 Positive Gaps and Strict Complementarity

In this section, we study the relationships between complementarity partitions and positive duality gaps. In particular, we consider cases where the complementarity partition for the pair of faces (f_P^0, f_D^0) fails to be strict. An instance with a finite positive gap is given in Example 2.5, item 2. We provide another example to illustrate the application of the optimality conditions that use the minimal sets.

Example 5.5 We let $K = K^* = \mathbb{S}_+^6$, with $\mathcal{A}^* y = \sum_i A_i y_i$ and

$$A_1 = E_{11}, A_2 = E_{22}, A_3 = E_{34}, A_4 = E_{13} + E_{55}, A_5 = E_{14} + E_{66}, \\ \tilde{y} = 0, c = \tilde{s} = E_{12} + E_{66}, b = (0 \ 0 \ 2 \ 0 \ 1)^T, \tilde{x} = E_{34} + E_{66}.$$

The primal and dual recession cones (first complementarity partition) are

$$f_P^0 = \begin{bmatrix} \mathbb{S}_+^2 & 0 \\ 0 & 0 \end{bmatrix} \trianglelefteq K, \quad f_D^0 = Q\mathbb{S}_+^2 Q^T \trianglelefteq K^*, \quad \text{where } Q = [e_3 \ e_4];$$

and positionwise

$$\begin{bmatrix} f_P^0 & 0 & 0 \\ 0 & f_D^0 & 0 \\ 0 & 0 & G \end{bmatrix},$$

where $G = 0_2$ represents the gap in strict complementarity. We note that $f_D^0 \cap \{c\}^\perp = f_D^0$. We apply (2.15) with the second (or fourth) complementarity partition in Proposition 2.3. We choose

$$x = \begin{bmatrix} 0_2 & 0 & 0 \\ 0 & I_2 & 0 \\ 0 & 0 & 0_2 \end{bmatrix} \in \text{relint } f_D^0 = \text{relint}(f_D^0 \cap \{c\}^\perp).$$

We get $f_P \triangleleft f_1 := K \cap \{x\}^\perp$ and the equivalent problem to (P)

$$(P) \quad v_P = \sup\{\langle b, y \rangle : \mathcal{A}^* y \preceq_{f_1} c\}.$$

However, we need one more step to find f_P . We again apply (2.15) and choose

$$0 \neq x = E_{55} \in \text{relint}(f_1 \cap \{c\}^\perp).$$

This yields $f_P = Q\mathbb{S}_+^3 Q^T$, where $Q = [e_1 \ e_2 \ e_6]$.

Similarly, we can work on the dual problem. In summary, we get that the faces and recession cones of the primal and dual are

$$f_P^0 = \begin{bmatrix} \mathbb{S}_+^2 & 0 \\ 0 & 0 \end{bmatrix} \trianglelefteq K, \quad f_D^0 = Q\mathbb{S}_+^2 Q^T \trianglelefteq K^*, \quad \text{where } Q = [e_3 \ e_4], \\ f_P = Q\mathbb{S}_+^3 Q^T, \quad \text{where } Q = [e_1 \ e_2 \ e_6], \quad f_D = Q\mathbb{S}_+^3 Q^T, \quad \text{where } Q = [e_3 \ e_4 \ e_6].$$

The optimal values are $v_P = 0$ and $v_D = 1$. ■

A connection between optimality and complementarity can be seen in the following proposition.

Proposition 5.6 Suppose that (2.2) has optimal solution \tilde{y} with corresponding optimal slack \tilde{s} , and that (2.3) has optimal solution \tilde{x} . Then

$$\langle \tilde{s}, \tilde{x} \rangle = \inf \{ \langle s, x \rangle : s \in \mathcal{F}_P^s, x \in \mathcal{F}_D^x \}.$$

Proof.

Since \tilde{s} is feasible, there exists $\ell' \in \mathcal{L}^\perp$ such that $c = \tilde{s} + \ell'$. Then for every feasible solution x of (2.3) and every feasible solution (y, s) of (2.2), we have

$$\begin{aligned} \langle s, x \rangle = \langle c, x \rangle - \langle b, y \rangle &= \langle \tilde{s} + \ell', x \rangle - \langle \mathcal{A}\tilde{x}, y \rangle \\ &= \langle \tilde{s}, x \rangle + \langle \ell', \tilde{x} \rangle - \langle \tilde{x}, \mathcal{A}^* y \rangle \\ &= \langle \tilde{s}, x \rangle - \langle \tilde{x}, c - s \rangle + \langle \ell', \tilde{x} \rangle \\ &= \langle \tilde{s}, x \rangle + \langle \tilde{x}, s \rangle + \underbrace{\langle \ell', \tilde{x} \rangle - \langle c, \tilde{x} \rangle}_{\text{constant}}. \end{aligned}$$

The second equation above uses the facts $\ell' \in \mathcal{L}^\perp$, $(\tilde{x} - x) \in \mathcal{L}$. Now, using the subspace form (2.9), (2.10) of the primal-dual pair, we conclude the desired result. \blacksquare

Simple necessary conditions for the existence of finite positive duality gap under the assumption that the complementarity partition (f_P^0, f_D^0) is strict, follow easily from Corollary 5.4.

Corollary 5.7 *Let K be a proper cone and suppose K and K^* are facially exposed. Suppose that \tilde{s} and \tilde{x} are optimal for (2.9) and (2.10), respectively, with a positive duality gap $\langle \tilde{s}, \tilde{x} \rangle > 0$. Moreover, suppose that the complementarity partition for the pair of faces (f_P^0, f_D^0) is strict. Then*

$$\tilde{x} \in \overline{(\text{face } \tilde{s})^c + (f_P^0)^\perp + \mathcal{L}} \setminus ((\text{face } \tilde{s})^c + \mathcal{L}), \quad \tilde{s} \in \overline{(\text{face } \tilde{x})^c + (f_D^0)^\perp + \mathcal{L}^\perp} \setminus ((\text{face } \tilde{x})^c + \mathcal{L}^\perp). \quad (5.5)$$

Proof.

The proof follows immediately from Corollary 5.4 after replacing $(f_D^0)^c$ and $(f_P^0)^c$ by f_P^0 and f_D^0 , respectively. \blacksquare

Example 5.8 (*S. Schurr [64]*) *It is possible to have a finite positive duality gap even if the complementarity partition for the pair of faces (f_P^0, f_D^0) is strict. Let $K = K^* = \mathbb{S}_+^5$, and*

$$\begin{aligned} A_1 = E_{11}, \quad A_2 = E_{22}, \quad A_3 = E_{34}, \quad A_4 = E_{13} + E_{45} + E_{55}, \\ b = (0 \quad 1 \quad 2 \quad 1)^T, \quad c = E_{44} + E_{55}. \end{aligned}$$

Then

$$f_P^0 = \begin{bmatrix} \mathbb{S}_+^2 & 0 \\ 0 & 0 \end{bmatrix}, \quad f_D^0 = \begin{bmatrix} 0 & 0 \\ 0 & \mathbb{S}_+^3 \end{bmatrix}, \quad f_P = Q\mathbb{S}_+^4Q^T, \quad f_D = \begin{bmatrix} 0 & 0 \\ 0 & \mathbb{S}_+^4 \end{bmatrix},$$

where $Q = [e_1 \ e_2 \ e_4 \ e_5]$. The primal optimal value is zero and the dual optimal value is $(\sqrt{5} - 1)/2$, and both are attained. This can be seen using the optimal $\tilde{s} = c$ for (2.2), and \tilde{x} optimal for (2.3) (optimal $x^* = \tilde{x}$ has values $1/\sqrt{5}$ and $(3 - \sqrt{5})/(2\sqrt{5})$ for the diagonal (5, 5) and (4, 4) elements, respectively). \blacksquare

We also have an example without the attainment of the optimal values.

Example 5.9 *Consider the SDP with data $K = K^* = \mathbb{S}_+^5$, and*

$$A_1 = E_{11}, \quad A_2 = E_{22}, \quad A_3 = E_{34}, \quad A_4 = E_{13} + E_{55}, \quad b = (0 \quad 1 \quad 2 \quad 1)^T, \quad c = E_{12} + E_{44} + E_{55}.$$

The primal optimal value is zero and the dual optimal value is 1, but neither value is attained. \blacksquare

We now consider cases when the assumption that the complementarity partition for the pair of faces (f_P^0, f_D^0) fails to be strict implies a finite positive duality gap. Our main result for the relationship between the failure of strict complementarity and finite nonzero duality gaps follows. We focus on the SDP case.

Theorem 5.10 *Let $K = \mathbb{S}_+^n$, suppose that the subspace $\mathcal{L} \subset \mathbb{S}^n$ is such that the complementarity partition for the pair of faces (f_P^0, f_D^0) minimally fails to be strict ($\dim G = 1$, where the face $G := (f_P^0)^c \cap (f_D^0)^c \leq K$). Then for every $M > 0$, there exist $\tilde{x}, \tilde{s} \in \text{relint } G$ such that the underlying problems (2.9) and (2.10) have a duality gap of exactly M .*

Proof.

First note that $\mathcal{L} \cap K = \{0\}$ if and only if $\mathcal{L}^\perp \cap \text{int } K \neq \emptyset$ if and only if $f_P^0 = K$. Using this and a similar result for $\mathcal{L}^\perp \cap K = \{0\}$, we conclude that both f_P^0 and f_D^0 are proper faces of K . Let $M > 0$ be arbitrary and $\tilde{s} \in \text{relint } G$ such that $\langle \tilde{s}, \tilde{s} \rangle = M$. In the definition of (\mathbb{D}) , let $\tilde{x} := \tilde{s}$. We claim that \tilde{s} is optimal in (\mathbb{P}) and (\mathbb{D}) . We prove the optimality claim by contradiction. Suppose \tilde{s} is not optimal in (\mathbb{P}) . Since it is feasible, with objective value M , there must exist another feasible solution of (\mathbb{D}) with strictly better objective value. The latter implies, there exists $u \in \mathbb{S}^n$ such that $u \in \mathcal{L}^\perp$, $u \succeq -\tilde{s}$ and $\langle \tilde{s}, u \rangle < 0$. Under an orthogonal similarity transformation, we have the following representation of the faces:

$$\begin{bmatrix} G & 0 & 0 \\ 0 & f_P^0 & 0 \\ 0 & 0 & f_D^0 \end{bmatrix}.$$

Then, $\tilde{s} = \sqrt{M}E_{11}$, and $u \succeq -\tilde{s}$ implies that the 3×3 block of u is zero. Thus,

$$u = \begin{bmatrix} \alpha & \bar{v}^T & 0 \\ \bar{v} & V & 0 \\ 0 & 0 & 0 \end{bmatrix},$$

where $\alpha < 0$. Now let $\hat{s} \in \text{relint } f_P^0$, that is,

$$\hat{s} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & \xi & 0 \\ 0 & 0 & 0 \end{bmatrix},$$

where $\xi > 0$. For every $\beta \in \mathbb{R}$, $(\beta\hat{s} - u) \in \mathcal{L}^\perp$. Moreover,

$$\beta\hat{s} - u = \begin{bmatrix} -\alpha & -\bar{v}^T & 0 \\ -\bar{v} & \beta\xi - V & 0 \\ 0 & 0 & 0 \end{bmatrix} \succ 0, \iff (\beta\xi - V) + \frac{1}{\alpha}\bar{v}\bar{v}^T \succ 0.$$

The latter is true for all sufficiently large $\beta > 0$. Hence, for all sufficiently large $\beta > 0$, $(\beta\hat{s} - u) \in \mathcal{L}^\perp \cap \mathbb{S}_+^n$ with $\text{rank}(\beta\hat{s} - u) = \text{rank}(\hat{s}) + 1$, a contradiction. This proves, \tilde{s} is optimal in (\mathbb{P}) . Similarly, \tilde{x} is optimal in (\mathbb{D}) . The duality gap is $\langle \tilde{x}, \tilde{s} \rangle = \langle \tilde{s}, \tilde{s} \rangle = M$, as claimed. \blacksquare

Example 5.11 *We now see that choosing one of \tilde{s}, \tilde{x} in $\text{relint } G$ may not result in a positive duality gap. Consider the SDP with data $K = K^* = \mathbb{S}_+^4$, and*

$$A_1 = E_{44}, \quad A_2 = E_{24} + E_{33}, \quad A_3 = E_{13} + E_{22}.$$

Then

$$f_P^0 = \begin{bmatrix} 0 & 0 \\ 0 & \mathbb{R}_+ \end{bmatrix} \trianglelefteq K, \quad f_D^0 = \begin{bmatrix} \mathbb{R}_+ & 0 \\ 0 & 0 \end{bmatrix} \trianglelefteq K^*,$$

and

$$G := (f_P^0)^c \cap (f_D^0)^c = QS_+^2 Q^T,$$

where $Q = [e_2 \ e_3]$. If \tilde{s} and \tilde{x} are chosen such that $\tilde{s} \in \text{relint}(G)$ and $\tilde{x} \in G$, with $\tilde{x}_{33} > 0$, then the optimal values are both $x_{33}(s_{33} - s_{23}^2/s_{22})$. However, there exist matrices $\tilde{s}, \tilde{x} \in G$ that are singular on G such that (2.2) and (2.3) admit a positive duality gap. For example, if $\tilde{s} = \tilde{x}$ is the diagonal matrix $\tilde{s} = \tilde{x} = \text{Diag}((0 \ 0 \ s_{33} \ 0))$, then the primal optimal value is zero and the dual optimal value is $x_{33}s_{33}$.¹ Both values are attained at $\tilde{s} = \tilde{x}$. To apply Corollary 5.4, we can write \tilde{x} as the limit

$$\begin{aligned} \tilde{x} &= x_{33} \lim_{i \rightarrow \infty} \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1/i \end{bmatrix} \\ &= x_{33} \lim_{i \rightarrow \infty} \begin{bmatrix} 0 & 0 & i & 0 \\ 0 & -2i & 0 & -1/2 \\ i & 0 & 1 & 0 \\ 0 & -1/2 & 0 & 0 \end{bmatrix} + \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 2i & 0 & 1/2 \\ 0 & 0 & 0 & 0 \\ 0 & 1/2 & 0 & 1/i \end{bmatrix} + \begin{bmatrix} 0 & 0 & -i & 0 \\ 0 & 0 & 0 & 0 \\ -i & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \\ &\in \overline{\mathcal{L} + (\text{face } \tilde{s})^c + ((f_D^0)^c)^\perp}. \end{aligned}$$

■

Note that the construction in the proof of Theorem 5.10 resulted in a primal-dual pair for which every feasible solution is optimal. We further investigate this connection in the next two results.

Theorem 5.12 *Let K be a closed convex cone. Suppose that the partition (f_P^0, f_D^0) is strictly complementary and that the following condition holds:*

$$\tilde{s} \in (f_D^0)^c + \mathcal{L}^\perp, \quad \tilde{x} \in (f_P^0)^c + \mathcal{L}. \quad (5.6)$$

Then every feasible solution in (P) and every feasible solution in (D) is optimal and strong duality holds for both (P) and (D).

Proof.

Suppose that (5.6) holds. Then by Proposition 3.4, part 2, $f_P \subseteq (f_D^0)^c$ and $f_D \subseteq (f_P^0)^c$. Further suppose that (f_P^0, f_D^0) forms a strict complementarity partition. Then $(f_D^0)^c = f_P^0$. Since for all feasible problems, $f_P^0 \subseteq f_P$ and $f_D^0 \subseteq f_D$ (Proposition 3.4, part 1), we actually have $f_P^0 = f_P$ and $f_D^0 = f_D$. Then, $\langle f_P, f_D \rangle = 0$, i.e. every feasible point is optimal (and we have no duality gap). ■

Corollary 5.13 *Suppose that both (2.9) and (2.10) are feasible but strong duality fails either problem. In addition, suppose that all feasible points for (2.2) and (2.3) are optimal. Then the complementarity partition for the pair of faces (f_P^0, f_D^0) fails to be strict.*

¹Similarly, we can use the 2, 2 position rather than the 3, 3 position.

Proof.

Suppose that all feasible points for (2.2) are optimal. Then the primal objective function is constant along all primal recession directions. That is, $\langle \tilde{x}, \mathcal{L}^\perp \cap K \rangle = \{0\}$, i.e., $\tilde{x} \in (\mathcal{L}^\perp \cap K)^\perp$. Now by construction, \tilde{x} is dual feasible, i.e., $\tilde{x} \in (\mathcal{L}^\perp \cap K)^\perp \cap K^* = (f_P^0)^\perp \cap K^* = (f_P^0)^c$. Finally, as argued previously, translating \tilde{x} by a point in \mathcal{L} leaves the dual problem unchanged, giving the condition on \tilde{x} in (5.6). In a similar way we can show that if all feasible points for (2.3) are optimal, then the condition on \tilde{s} in (5.6) holds. The desired result now follows from Theorem 5.12. \blacksquare

5.2 Infinite Duality Gap and Devious Faces

As we have already noted, (2.3) is feasible if and only if $\tilde{x} \in K^* + \mathcal{L}$. Moreover the feasibility of (2.3) is equivalent to a finite duality gap (possibly zero), recalling our assumption that the primal optimal value v_P is finite. We now see that if a nice cone has a devious face, then it is easy to construct examples with an infinite duality gap.

Proposition 5.14 *Suppose that K is a nice, proper cone and F is a devious face of K^* , i.e.,*

$$K^* + (F^c)^\perp = \overline{K^* + \text{span } F} \text{ and } (K^* + \text{span } F) \text{ is not closed.}$$

Let $\mathcal{L} = \text{span } F$ and choose $c = \tilde{s} = 0$ and $\tilde{x} \in (K^ + (F^c)^\perp) \setminus (K^* + \mathcal{L})$. Then $(\tilde{x} + \mathcal{L}) \cap K^* = \emptyset$ and we get $v_D = +\infty$. Moreover, $\mathcal{L}^\perp = F^\perp$ and, for every feasible $s \in F^\perp \cap K$,*

$$\langle \tilde{x}, s \rangle = \left\langle \tilde{x}_{K^*} + \tilde{x}_{(F^c)^\perp}, s \right\rangle \geq 0,$$

i.e., $0 = v_P < v_D = \infty$.

Proof.

The proof follows from the definitions. \blacksquare

Proposition 5.14 can be extended to choosing any \mathcal{L} that satisfies $K^* + \mathcal{L}$ is not closed and $K^* + \mathcal{L} \subset K^* + (F^c)^\perp$.

Example 5.15 *Let $K = \mathbb{S}_+^2$, and suppose that (2.2) and (2.3) admit a nonzero duality gap. Then Slater's CQ fails for both primal and dual, i.e., $\{0\} \neq f_P^0 \subset \mathbb{S}_+^2$ and $\{0\} \neq f_D^0 \subset \mathbb{S}_+^2$. After a rotation (see Lemma 2.8) we can assume the problem has the structure*

$$\begin{bmatrix} f_D^0 & 0 \\ 0 & f_P^0 \end{bmatrix},$$

viz., the matrices in f_D^0 are nonzero only in the (1,1) position, and the matrices in f_P^0 are nonzero only in the (3,3) position. There are only three possible options for \mathcal{L} : $\text{span}\{E_{11}\}$, $\text{span}\{E_{22}\}$, $\text{span}\{E_{11}, E_{12}\}$, or $\text{span}\{E_{22}, E_{12}\}$. In each case, either \mathcal{L} is one-dimensional and \mathcal{L}^\perp is two-dimensional, or vice versa. So without loss of generality, we may choose $\mathcal{L} = \text{span}\{E_{11}\}$. Now take $\tilde{x} = E_{12} \in \mathbb{S}_+^2 + (f_P^0)^\perp$. Then

$$\tilde{x} \notin \mathbb{S}_+^2 + \mathcal{L} = \mathbb{S}_+^2 + \text{span } f_D^0 \subset \mathbb{S}_+^2 + (f_P^0)^\perp, \quad (5.7)$$

and the dual program (2.3) is infeasible. But choosing $c = \tilde{s} = E_{22}$ implies that the primal optimal value $v_P = \langle c, \tilde{x} \rangle - y_1 \langle E_{22}, \tilde{x} \rangle = 0 < v_D = +\infty$.

Corollary 5.16 *If $K = \mathbb{S}_+^2$, then a finite positive duality gap cannot occur.*

Proof.

See Corollary 4.13. ■

The above Corollary 5.16 also follows from [66, Prop. 4], i.e., it states that a finite positive duality gap cannot happen if $\dim \mathcal{W} \leq 3$.

5.3 Regularization for Strict Complementarity

Suppose that strong duality holds for both the primal and dual SDPs, but strict complementarity fails for every primal-dual optimal solution $(\bar{s}, \bar{x}) \in \mathbb{S}_+^n \oplus \mathbb{S}_+^n$. Following [71], (\bar{s}, \bar{x}) is called a *maximal complementary solution pair* if the pair maximizes the sum $\text{rank}(s) + \text{rank}(x)$ over all primal-dual optimal (s, x) . The *strict complementarity nullity*, $g := n - \text{rank}(\bar{s}) - \text{rank}(\bar{x})$.

Let $\mathcal{U} = \mathcal{N}(\bar{s}) \cap \mathcal{N}(\bar{x})$ be the common nullspace of dimension g , and U be the $n \times g$ matrix with orthonormal columns satisfying $\mathcal{R}(U) = \mathcal{U}$. Let $[U \ Q]$ be an orthogonal matrix. Then we can regularize so that strict complementarity holds by replacing both primal-dual variables s, x by QsQ^T, QxQ^T , respectively. This is equivalent to replacing the matrices $C, A_i, i = 1, \dots, m$ that define \mathcal{A} by $Q^T C Q, Q^T A_i Q, i = 1, \dots, m$. (Note that we would then have to check for possible linear dependence of the new matrices $Q^T A_i Q$, as well as possible loss of Slater CQ.) Finding the common nullspace can be done dynamically during the solution process. This is done by checking the ratios of eigenvalues of s and x between iterates to see if the convergence is to 0 or to $O(1)$. (In the case of LP, this corresponds to identifying nonbasic variables using the so-called Tapia indices, see e.g., [30].)

6 Conclusion

In this paper we have looked at known and new, duality and optimality results for the cone optimization problem (2.2). We have used the subspace formulations of the primal and dual problems, (2.9),(2.10), to provide new CQs and new optimality conditions that hold without any CQ. This includes a UCQ, i.e., a CQ that holds independent of both data vectors b and c . In particular, the optimality characterizations show that a *minimal representation* of the cone and/or the linear transformation of the problem results in regularization, i.e., efficient modelling for the cone K and for the primal and dual constraints results in a stable formulation of the problem. In addition, we have discussed conditions for a zero duality gap and the surprising relations to the lack of strict complementarity in the homogeneous problem and to the closure of sums of cones. The (near) failure of Slater's CQ relates to both theoretical and numerical difficulties. The same holds true for the failure of strict complementarity. We have discussed regularization procedures for both failures. We hope that these results will lead to preprocessing for current cone optimization software packages.

¹ Research supported by The Natural Sciences and Engineering Research Council of Canada. Part of this paper is based on work presented at: ICCOPT II & MOPTA-07, McMaster University, Hamilton, 2007; 9th SIAM Conference on Optimization, Boston, May, 2008; and HPOPT 10, Tilburg, 2008. The authors would like to acknowledge Simon Schurr (Univ. of Waterloo) for many helpful conversations, as well as for other contributions to this paper. URL for paper: orion.math.uwaterloo.ca/~hwoikowi/henry/reports/ABSTRACTS.html

References

- [1] R.A. Abrams and A. Ben-Israel. Optimality conditions and recession cones. *Operations Res.*, 23(3):549–553, 1975. 6
- [2] F. Alizadeh and D. Goldfarb. Second-order cone programming. *Math. Program.*, 95(1, Ser. B):3–51, 2003. ISMP 2000, Part 3 (Atlanta, GA). 3
- [3] F. Alizadeh, J-P.A. Haeberly, and M.L. Overton. Primal-dual interior-point methods for semidefinite programming: Convergence rates, stability and numerical results. *SIAM J. Optim.*, 8:746–768, 1998. 18
- [4] E.D. Andersen, C. Roos, and T. Terlaky. Notes on duality in second order and p -order cone optimization. *Optimization*, 51(4):627–643, 2002. 3
- [5] M.F. Anjos and H. Wolkowicz. Strengthened semidefinite relaxations via a second lifting for the Max-Cut problem. *Discrete Appl. Math.*, 119(1-2):79–106, 2002. Foundations of heuristics in combinatorial optimization. 3
- [6] A. Auslender and M. Teboulle. *Asymptotic cones and functions in optimization and variational inequalities*. Springer Monographs in Mathematics. Springer-Verlag, New York, 2003. 6
- [7] G.P. Barker. The lattice of faces of a finite dimensional cone. *Linear Algebra and Appl.*, 7:71–82, 1973. 8
- [8] G.P. Barker. Faces and duality in convex cones. *Linear and Multilinear Algebra*, 6(3):161–169, 1978/79. 8
- [9] A. Ben-Israel, A. Ben-Tal, and S. Zlobec. *Optimality in nonlinear programming: a feasible directions approach*. John Wiley & Sons Inc., New York, 1981. A Wiley-Interscience Publication. 7
- [10] A. Berman. *Cones, Matrices and Mathematical Programming*. Springer-Verlag, Berlin, New York, 1973. 17, 20
- [11] D.P. Bertsekas and P. Tseng. Set intersection theorems and existence of optimal solutions. *Math. Program.*, 110(2, Ser. A):287–314, 2007. 6
- [12] J.M. Borwein and W.B. Moors. Stability of closedness of convex cones under linear mappings. Technical report, Dalhousie University, Halifax, N.S., 2008. 17
- [13] J.M. Borwein and H. Wolkowicz. Characterization of optimality for the abstract convex program with finite-dimensional range. *J. Austral. Math. Soc. Ser. A*, 30(4):390–411, 1980/81. 3, 12, 14, 15
- [14] J.M. Borwein and H. Wolkowicz. Facial reduction for a cone-convex programming problem. *J. Austral. Math. Soc. Ser. A*, 30(3):369–380, 1980/81. 3, 14
- [15] J.M. Borwein and H. Wolkowicz. Regularizing the abstract convex program. *J. Math. Anal. Appl.*, 83(2):495–530, 1981. 3, 13, 14

- [16] J.M. Borwein and H. Wolkowicz. Characterizations of optimality without constraint qualification for the abstract convex program. *Math. Programming Stud.*, 19:77–100, 1982. Optimality and stability in mathematical programming. 3
- [17] A. Brøndsted. Les faces d’un cône convexe abstrait. *C. R. Acad. Sci. Paris Sér. A-B*, 280(7):Aii, A435–A437, 1975. 8
- [18] Y-L. Cheung, S. Schurr, and H. Wolkowicz. Numerical solution of ill-posed semidefinite programs. Technical Report CORR 2010, University of Waterloo, Waterloo, Ontario, 2010. in progress. 3, 16
- [19] C.B. Chua. Relating homogeneous cones and positive definite cones via T -algebras. *SIAM J. Optim.*, 14(2):500–506 (electronic), 2003. 13
- [20] C.B. Chua and L. Tunçel. Invariance and efficiency of convex representations. *Math. Program.*, 111(1-2, Ser. B):113–140, 2008. 13
- [21] F.H. Clarke. *Optimization and Nonsmooth Analysis*. Canadian Math. Soc. Series of Monographs and Advanced Texts. John Wiley & Sons, 1983. 7
- [22] F. Deutsch. The role of the strong conical hull intersection property in convex optimization and approximation. In *Approximation theory IX, Vol. I. (Nashville, TN, 1998)*, Innov. Appl. Math., pages 105–112. Vanderbilt Univ. Press, Nashville, TN, 1998. 17
- [23] R.J. Duffin. Infinite programs. In A.W. Tucker, editor, *Linear Equalities and Related Systems*, pages 157–170. Princeton University Press, Princeton, NJ, 1956. 3
- [24] L. Faybusovich. On Nesterov’s approach to semi-infinite programming. *Acta Appl. Math.*, 74(2):195–215, 2002. 13
- [25] R.M. Freund, F. Ordóñez, and K.C. Toh. Behavioral measures and their correlation with IPM iteration counts on semi-definite programming problems. USC-ISE working paper #2005-02, MIT, 2005. url: www-rcf.usc.edu/~fordon/. 3, 18
- [26] R.M. Freund and J.R. Vera. Condition-based complexity of convex optimization in conic linear form via the ellipsoid algorithm. *SIAM J. Optim.*, 10(1):155–176 (electronic), 1999. 3, 18
- [27] K. Fujisawa, M. Fukuda, and K. Nakata. Preprocessing sparse semidefinite programs via matrix completion. *Optim. Methods Softw.*, 21(1):17–39, 2006. 3
- [28] A.J. Goldman and A.W. Tucker. Theory of linear programming. In *Linear inequalities and related systems*, pages 53–97. Princeton University Press, Princeton, N.J., 1956. Annals of Mathematics Studies, no. 38. 3
- [29] J. Gondzio. Presolve analysis of linear programs prior to applying an interior point method. *INFORMS J. Comput.*, 9(1):73–91, 1997. 3
- [30] M. Gonzalez-Lima, H. Wei, and H. Wolkowicz. A stable primal-dual approach for linear programming under nondegeneracy assumptions. *Comput. Optim. Appl.*, 44(2):213–247, 2009. 26

- [31] M. Guignard. Generalized Kuhn-Tucker conditions for mathematical programming problems in a Banach space. *SIAM J. of Control*, 7(2):232–241, 1969. 3, 16, 17
- [32] M. Halická, E. de Klerk, and C. Roos. On the convergence of the central path in semidefinite optimization. *SIAM J. Optim.*, 12(4):1090–1099 (electronic), 2002. 18
- [33] C. Helmberg. Semidefinite programming. *European J. Oper. Res.*, 137(3):461–482, 2002. 3
- [34] R.B. Holmes. *Geometric Functional Analysis and its Applications*. Springer-Verlag, Berlin, 1975. 17, 20
- [35] V. Jeyakumar and G.M. Lee. Complete characterizations of stable Farkas’ lemma and cone-convex programming duality. *Math. Program.*, 114(2, Ser. A):335–347, 2008. 3
- [36] V. Jeyakumar and G.Y. Li. Stable zero duality gaps in convex programming: complete dual characterisations with applications to semidefinite programs. *J. Math. Anal. Appl.*, 360(1):156–167, 2009. 3
- [37] J. Ji, F.A. Potra, and R. Sheng. On the local convergence of a predictor-corrector method for semidefinite programming. *SIAM J. Optim.*, 10(1):195–210 (electronic), 1999. 18
- [38] D. Jibetean and M. Laurent. Semidefinite approximations for global unconstrained polynomial optimization. *SIAM J. Optim.*, 16(2):490–514 (electronic), 2005. 3
- [39] A. Koberstein and U.H. Suhl. Progress in the dual simplex method for large scale LP problems: practical dual phase 1 algorithms. *Comput. Optim. Appl.*, 37(1):49–65, 2007. 3
- [40] M. Kojima, M. Shida, and S. Shindoh. Local convergence of predictor-corrector infeasible-interior-point algorithms for SDPs and SDLCPs. Technical report, Dept. of Information Sciences, Tokyo Institute of Technology, Tokyo, Japan, 1996. 18
- [41] K. Krishnan and T. Terlaky. Interior point and semidefinite approaches in combinatorial optimization. In *Graph theory and combinatorial optimization*, volume 8 of *GERAD 25th Anniv. Ser.*, pages 101–157. Springer, New York, 2005. 3
- [42] N. Krislock and H. Wolkowicz. Explicit sensor network localization using semidefinite representations and facial reductions. *SIAM Journal on Optimization*, 20(5):2679–2708, 2010. 3
- [43] S. Kurcyusz. On the existence of Lagrange multipliers for infinite dimensional extremal problems. *Bull. Acad. Polon. Sci. Sér. Sci. Tech.*, 22:799–802, 1974. 17
- [44] S. Kurcyusz. On the existence and non-existence Lagrange multipliers in Banach spaces. *J. Optimization Theory Appl.*, 20(1):81–110, 1976. 17
- [45] M.S. Lobo, L. Vandenberghe, S. Boyd, and H. Lebret. Applications of second-order cone programming. *Linear Algebra Appl.*, 284(1-3):193–228, 1998. ILAS Symposium on Fast Algorithms for Control, Signals and Image Processing (Winnipeg, MB, 1997). 3
- [46] D.G. Luenberger. *Optimization by Vector Space Methods*. John Wiley, 1969. 14, 19

- [47] Z.-Q. Luo, J. F. Sturm, and S. Zhang. Superlinear convergence of a symmetric primal-dual path-following algorithm for semidefinite programming. *SIAM J. Optim.*, 8:59–81, 1998. 18
- [48] C. Mészáros and J. Gondzio. Addendum to: “Presolve analysis of linear programs prior to applying an interior point method” [INFORMS J. Comput. **9** (1997), no. 1, 73–91; MR 99a:90120] by Gondzio. *INFORMS J. Comput.*, 13(2):169–170, 2001. 3
- [49] R.D.C. Monteiro. First- and second-order methods for semidefinite programming. *Math. Program.*, 97(1-2, Ser. B):209–244, 2003. ISMP, 2003 (Copenhagen). 3
- [50] Y.E. Nesterov and A.S. Nemirovski. *Interior Point Polynomial Algorithms in Convex Programming*. SIAM Publications. SIAM, Philadelphia, USA, 1994. 5
- [51] A.G. Pais. Cone extreme points and faces. *J. Math. Anal. Appl.*, 126(1):223–228, 1987. 8
- [52] G. Pataki. On the closedness of the linear image of a closed convex cone. *Math. Oper. Res.*, 32(2):395–412, 2007. 12, 13, 17
- [53] I. Pólik and T. Terlaky. Exact duality for optimization over symmetric cones. Technical report, McMaster University, Hamilton, Ontario, Canada, 2007. 3, 13
- [54] F.A. Potra and R. Sheng. Superlinear convergence of a predictor-corrector method for semidefinite programming without shrinking central path neighborhood. *Bull. Math. Soc. Sci. Math. Roumanie (N.S.)*, 43(91)(2):107–124, 2000. 18
- [55] B.N. Pshenichniyi. *Necessary conditions for an extremum*, volume 4 of *Translated from the Russian by Karol Makowski. Translation edited by Lucien W. Neustadt. Pure and Applied Mathematics*. Marcel Dekker Inc., New York, 1971. 17
- [56] M.V. Ramana. *An Algorithmic Analysis of Multiquadratic and Semidefinite Programming Problems*. PhD thesis, Johns Hopkins University, Baltimore, Md, 1993. 14
- [57] M.V. Ramana. An exact duality theory for semidefinite programming and its complexity implications. *Math. Programming*, 77(2):129–162, 1997. 3, 13, 14
- [58] M.V. Ramana, L. Tunçel, and H. Wolkowicz. Strong duality for semidefinite programming. *SIAM J. Optim.*, 7(3):641–662, 1997. 3, 12, 13, 14
- [59] J. Renegar. Some perturbation theory for linear programming. *Math. Programming*, 65(1, Ser. A):73–91, 1994. 18
- [60] J. Renegar. Incorporating condition measures into the complexity theory of linear programming. *SIAM J. Optim.*, 5(3):506–524, 1995. 3
- [61] J. Renegar. Linear programming, complexity theory and elementary functional analysis. *Math. Programming*, 70(3, Ser. A):279–351, 1995. 3, 18
- [62] S. M. Robinson. Stability theorems for systems of inequalities, part ii: differentiable nonlinear systems. *SIAM J. Numerical Analysis*, 13:497–513, 1976. 3
- [63] R.T. Rockafellar. *Convex analysis*. Princeton Landmarks in Mathematics. Princeton University Press, Princeton, NJ, 1997. Reprint of the 1970 original, Princeton Paperbacks. 10, 11, 13

- [64] S.P. Schurr. Private communication. Technical report, University of Waterloo, Waterloo, ON, 2008. 22
- [65] A. Shapiro. Duality and optimality conditions. In *Handbook OF Semidefinite Programming: Theory, Algorithms, and Applications*. Kluwer Academic Publishers, Boston, MA, 2000. 2, 3, 14
- [66] A. Shapiro and A. Nemirovskii. Duality of linear conic problems. Technical report, School of Industrial and Systems Engineering, Georgia Institute of Technology, Atlanta, Georgia, 2003. 7, 13, 26
- [67] M.J. Todd. Semidefinite programming. *Acta Numerica*, 10:515–560, 2001. 3
- [68] L. Tunçel. On the Slater condition for the SDP relaxations of nonconvex sets. *Oper. Res. Lett.*, 29(4):181–186, 2001. 3
- [69] H. Waki, S. Kim, M. Kojima, and M. Muramatsu. Sums of squares and semidefinite program relaxations for polynomial optimization problems with structured sparsity. *SIAM J. Optim.*, 17(1):218–242, 2006. 3
- [70] Z. Waksman and M. Epelman. On point classification in convex sets. *Math. Scand.*, 38(1):83–96, 1976. 7
- [71] H. Wei and H. Wolkowicz. Generating and solving hard instances in semidefinite programming. *Math. Programming*, 125(1):31–45, 2010. 3, 8, 18, 26
- [72] H. Wolkowicz. Some applications of optimization in matrix theory. *Linear Algebra Appl.*, 40:101–118, 1981. 14
- [73] H. Wolkowicz, R. Saigal, and L. Vandenberghe, editors. *Handbook of semidefinite programming*. International Series in Operations Research & Management Science, 27. Kluwer Academic Publishers, Boston, MA, 2000. Theory, algorithms, and applications. 3, 12, 13, 14
- [74] H. Wolkowicz and Q. Zhao. Semidefinite programming relaxations for the graph partitioning problem. *Discrete Appl. Math.*, 96/97:461–479, 1999. Selected for the special Editors’ Choice, Edition 1999. 3
- [75] Q. Zhao, S.E. Karisch, F. Rendl, and H. Wolkowicz. Semidefinite programming relaxations for the quadratic assignment problem. *J. Comb. Optim.*, 2(1):71–109, 1998. Semidefinite programming and interior-point approaches for combinatorial optimization problems (Fields Institute, Toronto, ON, 1996). 3
- [76] J. Zowe and S. Kurcyusz. Regularity and stability for the mathematical programming problem in Banach spaces. *Appl. Math. Optim.*, 5:49–62, 1979. 17
- [77] L.F. Zuluaga, J. Vera, and J. Peña. LMI approximations for cones of positive semidefinite forms. *SIAM J. Optim.*, 16(4):1076–1091 (electronic), 2006. 3

Index

- $\mathcal{A} : \mathcal{V} \rightarrow \mathcal{W}$, linear transformation, 4
 $\mathcal{F}_P^s = \mathcal{F}_P^s(c)$, primal feasible slacks, 5
 $\mathcal{N}(\cdot)$, nullspace, 4
 $\mathcal{O}_P^s, \mathcal{O}_P^y, \mathcal{O}_D^x$, optimal solutions sets, 5
 $\mathcal{R}(\cdot)$, range space, 4, 16
 $\text{precl}S$, preclosure of S , 4
 $u \preceq_K v$, partial order induced by K , 4
 FDC, facially dual-complete (nice) cone, 12
 $\mathcal{D}_P^{\leq}(\hat{y}), \mathcal{D}_P^{\leq}(\hat{s}), \mathcal{D}_D^{\leq}(\hat{x})$, cones of feasible directions, 7
 closure of S , \bar{S} , 4
 complementarity partition, CP, of K, K^* , 6
 cones of feasible directions, $\mathcal{D}_P^{\leq}(\hat{y}), \mathcal{D}_P^{\leq}(\hat{s}), \mathcal{D}_D^{\leq}(\hat{x})$, 7
 conjugate face, 4
 convex cone generated by S , cone S , 4
 convex hull, $\text{conv } S$, 4
 CP, complementarity partition of K, K^* , 6
 devious cone, 13
 devious face, $K^* + \text{span } F$ not closed, 13
 dual feasible set, $\mathcal{F}_D^x = \mathcal{F}_D^x(b)$, 5
 dual pair of regularized programs based on primal minimal face, 14
 dual pair of regularized programs based on primal minimal subspace representation, $(\text{RP}_M), (\text{DRP}_M)$, 15
 exposed face, 4, 9
 face, $F \trianglelefteq K$, 4
 facially exposed cone, 4
 generalized Slater CQ, 16
 maximal complementary solution pair, 26
 minimal faces, $f_P = f_P(\tilde{s}), f_D = f_D(\tilde{x})$, 6
 minimal subspace representation, \mathcal{L}_{PM}^\perp , 15
 minimal subspace representations, $\mathcal{L}_{PM}^\perp, \mathcal{L}_{DM}$, 7
 nice cone, 12
 nice face, FDC, $K^* + F^\perp$ is closed, 12
 nullspace, $\mathcal{N}(\cdot)$, 4
 optimal solutions sets, $\mathcal{O}_P^s, \mathcal{O}_P^y, \mathcal{O}_D^x$, 5
 pair of dual conic optimization problems, 4
 partial order induced by K , $u \preceq_K v$, 4
 preclosure of S , $\text{precl}S$, 4
 primal feasible sets, $\mathcal{F}_P^y = \mathcal{F}_P^y(c)$, 5
 primal feasible slacks, $\mathcal{F}_P^s = \mathcal{F}_P^s(c)$, 5
 proper complementarity partition of K, K^* , 6
 proper cone, 4
 proper face, 4
 proper subset, 4
 range space, $\mathcal{R}(\cdot)$, 4
 Rockafellar-Pshenichnyi optimality condition, 17, 19
 smallest face containing a set $S \subseteq K$, $\text{face}(S) = \text{face } S$, 9
 strict complementarity, 5
 strict complementarity nullity, 26
 strict complementarity partition of K, K^* , 6
 strong duality, 5, 14
 subspace CQ, $\mathcal{R}(\mathcal{A}^*) \subseteq f_P - f_P$, 16
 subspace form for (2.2),(2.3), 5
 support of $x \in \mathbb{S}^n$, $\mathcal{S}(x)$, 8
 tangent cones, $\mathbb{T}_P(\hat{y}), \mathbb{T}_P(\hat{s}), \mathbb{T}_D(\hat{x})$, 7
 trace inner-product, trace sx , 4
 universal CQ, UCQ, 14, 18
 weak duality, 5
 weakest constraint qualification, WCQ, 14

A comments from referee

Question from referee:

Thm 5.3 seems to be incorrect and in this case Cor 5.4, Cor 5.7 would be incorrect as well. Am I missing something? Example 4 from “Ramana, An exact duality theory...” with $\alpha = 1$ seems to contradict Thm 5.3:

1. $A_1 = E_{22}, A_2 = E_{11} + E_{23}, C = E_{11}, \tilde{s} = E_{11} + E_{22}, \tilde{x} = E_{11}$.
2. $\langle \tilde{s}, \tilde{x} \rangle > 0$.
3. $(K - \tilde{s})^*$ is the cone generated by E_{33} .
4. \mathcal{L} is the subspace of symmetric matrices orthogonal to A_1 and A_2 .
5. Therefore, $(K - \tilde{s})^* \subseteq \mathcal{L}$ and the preclosure of $(K - \tilde{s})^* + \mathcal{L}$ is empty.